



Durham E-Theses

Late Jurassic Sedimentation in the Boreal-Tethyan Seaway: climate modelling, geochemistry, and petrography of the Kimmeridge Clay Formation

ATAR, ELIZABETH,FILIZ,LINDSAY

How to cite:

ATAR, ELIZABETH,FILIZ,LINDSAY (2019) *Late Jurassic Sedimentation in the Boreal-Tethyan Seaway: climate modelling, geochemistry, and petrography of the Kimmeridge Clay Formation*, Durham theses, Durham University. Available at Durham E-Theses Online: <http://etheses.dur.ac.uk/13132/>

Use policy

The full-text may be used and/or reproduced, and given to third parties in any format or medium, without prior permission or charge, for personal research or study, educational, or not-for-profit purposes provided that:

- a full bibliographic reference is made to the original source
- a [link](#) is made to the metadata record in Durham E-Theses
- the full-text is not changed in any way

The full-text must not be sold in any format or medium without the formal permission of the copyright holders.

Please consult the [full Durham E-Theses policy](#) for further details.

Appendix A - Climate modelling figures




Figure 0.1. Preindustrial simulation surface temperature maps. (a, d, and g) Annual, winter (DJF), and summer (JJA) surface temperature maps from HadCM3L. (b, e, h) Annual, winter (DJF), and summer (JJA) surface temperature maps from FOAM. (c, f, and i) Annual, winter (DJF), and summer (JJA) temperature differences between HadCM3L and FOAM.

Appendix A




Figure 0.2. Preindustrial simulation precipitation maps. (a, d, and g) Annual, winter (DJF), and summer (JJA) precipitation from HadCM3L. (b, e, h) Annual, winter (DJF), and summer (JJA) precipitation from FOAM. (c, f, and i) Annual, winter (DJF), and summer (JJA) precipitation differences between HadCM3L and FOAM.

Near Surface temperature (°C)	HadCM3L	FOAM	HadCM3L/FOAM Difference
Annual	14.6	7.3	7.3
DJF	12.3	5.1	7.2
JJA	16.5	8.8	7.7

Table 0.1. Preindustrial annual, winter (DJF), and summer (JJA) global mean surface temperatures for the HadCM3L and FOAM and the difference between the two models.




Figure 0.3. Boreal summer (JJA) and winter (DJF) mean precipitation (mm/s) from HadCM3L (left) and FOAM (right). Overlain, the red lines show the local maximum in vertical atmospheric ascent velocity (w) at a height of 500 mbar, for regions equator-wards of 30 degrees N/S, with 500 mbar temperature of greater than 260 K, and $w > 0.005$ m/s. The blue lines show the local maximum in precipitation (p), for regions equator-wards of 30 degrees N/S, with 500 mbar temperature of greater than 260K, and $p > 0.00003$ mm/s. Thick black line encompasses dry regions where the precipitation rate is less than $1 \cdot 10^{-5}$ mm/s. The orange line encompasses regions of mid-atmospheric descent where the vertical velocity at 500 mbar is towards the surface and greater than 0.02 is m/s. After Armstrong et al., (2016).