

Durham E-Theses

Quaternary sea level change, Lake formation, and associated glacial events, with special reference to the lower tees basin

Lockery, Andrew R.

How to cite:

Lockery, Andrew R. (1971) *Quaternary sea level change, Lake formation, and associated glacial events, with special reference to the lower tees basin*, Durham theses, Durham University. Available at Durham E-Theses Online: <http://etheses.dur.ac.uk/8445/>

Use policy


The full-text may be used and/or reproduced, and given to third parties in any format or medium, without prior permission or charge, for personal research or study, educational, or not-for-profit purposes provided that:

- a full bibliographic reference is made to the original source
- a [link](#) is made to the metadata record in Durham E-Theses
- the full-text is not changed in any way

The full-text must not be sold in any format or medium without the formal permission of the copyright holders.

Please consult the [full Durham E-Theses policy](#) for further details.

ISOSTATIC UPLIFT




ADAPTED FROM NANSSEN 1927

FIG. 1.




THE EFFECTS OF DIFFERENT RATES OF
GLACIER RETREAT ON THE AMOUNT OF
POST GLACIAL REBOUND



AFTER ANDREWS 1968

FIG. 2


ISOBASES OF LATE AND POSTGLACIAL RAISED SHORELINES IN
HIGHLAND BRITAIN




AFTER SMITH

FIG. 3


GRAPHS DEPICTING THE EUSTATIC RISE OF SEA LEVEL FOLLOWING THE LAST ICE AGE




1.




3.



2.



4.




5.

- 1) Fairbridge 1961
- 2) Godwin and Suggate 1958
- 3) Shepard and Suess 1956
- 4) Curray 1960
- 5) Shepard 1960


FIG. 4

GRAPHS DEPICTING THE EUSTATIC RISE OF SEA LEVEL FOLLOWING THE LAST ICE AGE




Sea level change in the
Fenlands of East Anglia

(Godwin 1940)



(New Zealand)

(Schofield 1960)



10000 BP
(Dutch Coast)

5000 BP

(Jelgersma 1961)

FIG. 5

RETREAT OF THE WÜRM ICE SHEET



D DISTRICT
 --- LAMMERMUIR STRANRAAR --- HIGHLAND
 — N.E. ANTRIM I.O.M. CUMBERLAND
 — NEWDRIFT
Mc CALLIEN
MOVIUS

3 DINNET RE-ADVANCE
 2 ABERDEEN RE-ADVANCE
 1 MORAY FIRTH
 --- STRATHMORE GLACIATION
 --- OTHER STAGES
SYNGE



..... MORaine GLACIATION
 --- HIGHLAND
 --- NORTHERN BRITISH RE-ADVANCE
CHARLESWORTH

AFTER SMITH



— LOCH LOMOND
 --- PERTH
 --- LAMMERMUIR
 --- YORK RE-ADVANCE
SISSONS

FIG. 6





FIG. 7

THE LOWER TEES.






(BELOW 100)

Relief and Drainage.

FIG. 8

THE LOWER TEES.

(Below 100')



Amplitude of Relief.

THE LOWER TEES.

(BELOW 100.)





FIG.10
Areas unsuitable for morphometric analysis. BASED ON PELLONCK, D.S. STAMP, AND TESSIER, CHART 1901.


THE LOWER TEES.



Drift Geology.

FIG. 11


GLACIAL RETREAT STAGES IN NORTH-WEST ENGLAND




FROM A. RAISTRICK

Fig. 12

THE LOWER TEES.




PROFILES CONSTRUCTED FROM
CONTOURS DRAWN AT 10 INTERVALS
USING AVAILABLE SPOT HEIGHTS
AND BENCH MARKS.



0 1 mile 2
SCALE.

Location of topographic profiles.

**PROFILES ACROSS 10 FOOT CONTOUR INTERVALS, CONSTRUCTED
FROM SPOT HEIGHTS AND BENCH MARKS.**



0 ————— 1
MILE

(NUMBERS REFER TO THOSE ON FIG. 13)

FIG-14

THE LOWER TEES
(BELOW 100 FEET)
LOCATION OF LEVELLED
TRAVERSES






FIG. 15

RANGE 2



HARTLEPOOL

TEES BAY




1 MILE

ECHO-SOUNDING

DIRECTION	North East
SPEED	1.5 Knots
WIND SPEED	Force 5
CONDITIONS	16 ft swell
DATE	6 12 67 11c

THE LOWER TEES BASIN
 HEIGHTS OF LEVELLED TRAVERSES SOUTH OF THE R. TEES
 — STAINSBY HALL FARM TO SALT BURN



THE LOWER TEES BASIN
 HEIGHTS OF LEVELLED TRAVERSES SOUTH OF THE R. TEES
 STAINSBY HALL FARM TO LOW HAIL FARM



FIG. 17

THE LOWER TEES BASIN
HEIGHTS OF LEVELLED TRAVERSES NORTH OF THE R. TEES
— HURWORTH TO BILLINGHAM BECK



THE LOWER TEES BASIN
 HEIGHTS OF LEVELLED TRAVERSES NORTH OF THE R. TEES
 — WOLVISTON TO HARTLEPOOL




FIG. 19

THE LOCATION OF ECHO SOUNDINGS OFF THE EAST COAST OF HIGHLAND
BRITAIN SHOWING THE FIVE MAJOR REGIONS





FIG. 21

THE TEES BAY
THE LOCATION OF SOUNDINGS

GROUP I





FIG. 22

TEES BAY TO TYNEMOUTH
LOCATION OF SOUNDINGS





0 5
MILES

----- SELF
----- COMMERCIAL

FIG. 23

TYNEMOUTH TO BERWICK
THE LOCATION OF SOUNDINGS



0 5
MILES

FIG. 24

ABBS HEAD AND THE FIRTH OF FORTHTHE LOCATION OF
SOUNDINGS

FIRTH OF FORTH

ABBS HEAD

BERWICK




FIG. 25

FIRTH OF FORTH TO STONEHAVEN
THE LOCATION OF SOUNDINGS




FIG. 26




FIG. 27

SEABED PROFILES
THE LOWER TEES

REBCAR

- 113'

S. GARE
LIGHTHOUSE

- 70'

- 80'

SLIGHT CHANGE
OF GRADIENT

- 180'

SLIGHT
CHANGE
OF GRADI-
- ENT

N. GARE
LIGHTHOUSE

- 70'

- 80'

- 178'

LONG SCAR ROCKS

- 70'

- 80'

HARTLEPOOL (SOUTH)

- 180'

HARTLEPOOL (NORTH)

- 180'



NAUTICAL MILE

VERTICAL SCALE : X 2

FIG. 28

SEABED CONTOUR PLAN OFF THE SUNDERLAND COAST

SOUNDINGS ARE SHOWN IN FEET
REDUCED TO ADMIRALTY CHART
DATUM.

CONTOURS AT 5FT. INTERVALS.

54° 53' 00"


54° 53' 00"

54° 51' 00"

54° 51' 00"

SEAHAM

54° 49' 00"




1000 0 10000
SCALE IN FT.

1000 0 5000
SCALE IN METERS

FIG. 29

54° 47' 00"



**SOUNDINGS COMPLETED BY
LAND & MARINE CONTRACTORS
LTD. IMMEDIATELY SOUTH OF
TYNEMOUTH, ENGLAND**



SOUNDINGS IN FEET

ALL SOUNDINGS REDUCED
TO CHART DATUM.

CHART DATUM IS 6.88 FT.
BELOW ORDNANCE DATUM
(NEWLYN) I.E. O.D. - 6.88 FT.

CONTOURS 0' TO 50' AT 5' INTERVALS

CONTOURS 50' TO 120' AT 2' INTERVALS



SCALE IN FT.

FIG. 30

**SOUNDINGS COMPLETED BY LAND & MARINE CONTRACTORS
LTD. IMMEDIATELY NORTH OF WHITVEY BAY**

H.W.M.M.T.

ST. MARY'S LIGHTHOUSE

SOUNDINGS IN FEET
ALL SOUNDINGS REDUCED
TO CHART DATUM.

CHART DATUM IS 6.88 FT.
BELOW ORDNANCE DATUM
(NEWLYN) I.E. O.D. - 6.88 FT.

CONTOURS 0' TO 50' AT 5' INTERVALS

CONTOURS 50' TO 120 AT 2' INTERVALS

0 1000 2000 3000 4000 5000

SCALE IN FT.




FIG. 31

SEABED CONTOURS OFF THE SOUTHERN NORTHUMBERLAND COAST

BASED ON SOUNDINGS
COMPLETED BY THE
DOVE MARINE LABORATORY

CONTOURS & FIGURES
SHOW DEPTH IN METERS
(FEET) IN BRACKETS
BELOW CHART DATUM.

CHART DATUM APPROX.
7-FT. BELOW O.D.




FIG. 32

SOUTER POINT (TYNEMOUTH)

SEABED PROFILES

BLYTHE
(NORTHUMBERLAND)

ST. ABBS HEAD

FIRTH OF FORTH

ST. ANDREW'S BAY

ARBROATH TO STONEHAVEN



NAUTICAL MILE

VERTICAL SCALE:X2

FIG. 33

SCHEMATIC DIAGRAM OF SPARKER SURVEY
AND EQUIPMENT





FIG. 34

PROFILE I

E.N.E FROM POINT ON SHORE MIDWAY BETWEEN MARSKE AND REDCAR





HORIZONTAL SCALE : 1" = 1700'
VERTICAL SCALE : 1" = 170'
VERTICAL EXAGGERATION: X10

NOTE: PROFILE SHOWS ONLY BEDROCK

FIG. 35

E.N.E. FROM POINT MIDWAY BETWEEN WEST SCAR ROCKS & OFF THE SOUTH GARE LIGHTHOUSE

OD. NEWLYN




HORIZONTAL SCALE: 1" = 1700'

VERTICAL SCALE: 1" = 170'

VERTICAL EXAGGERATION: X10

FIG. 36

EXPERIMENT ILLUSTRATING THE POSSIBLE RELATIONSHIP BETWEEN SEABED FORMS AND
THE SURFACE OF THE UNDERLYING BOULDER CLAY.



NOTE HOW THE SETTLED LAYERS HAVE THE EFFECT
OF MOVING THE BREAK OF SLOPE TOWARD DEEPER WATER

PROFILE 3

E.N.E FROM POINT ONE MILE SOUTH OF THE SOUTH GARE LIGHTHOUSE




FIG 38

N.N.E. FROM NORTH GARE LIGHTHOUSE




FIG. 39

GRAPHICAL METHODS OF PRESENTATION OF PARTICLE SIZE ANALYSES









FIG. 40.

THE SEPARATION OF BEACH AND DUNE SANDS


USING FRIEDMAN'S MEASURE OF SKEW




After Chappell 1967

FIG. 41

SAMPLING DIAGRAMS
ILLINOIS BEACH EXPERIMENT




A. SIMPLE RANDOM SAMPLES




B. RANDOM IN CELLS

→ N




C. SYSTEMATIC IN CELLS




D. CLUSTERS OF FOUR

AFTER KRUMBEIN & SLACK


SAMPLING DIAGRAMS
ILLINOIS BEACH EXPERIMENT




E. THREE LEVEL DESIGN



F. STRATIFIED RANDOM



G. THREE STRATA (COMBINED SAMPLES)




H. PURPOSIVE SELECTION

AFTER KRUMBEIN & SLACK

FIG. 43

**CONFIDENCE BANDS AND RELATIVE EFFICIENCY OF SAMPLING
DESIGNS FOR PHI MEAN PARTICLE DIAMETERS.**



AFTER KRUMBEIN & SLACK

FIG. 44.

STRATIFIED RANDOM SAMPLES ON THE PRESENT DAY BEACH AT REDCAR







FIG. 4.5

MEAN CURVES OF SAND SIZE FRACTIONS




FIG. 46

MAXIMUM VARIATIONS EXHIBITED BY PARTICLE SIZE CURVES







FIG. 47

THORPE THEWLES BECK
THE EFFECTS OF CHANGING CONDITION ON PARTICLE SIZE




FIG. 48

THE LOCATION OF BEACH SAMPLE AREAS
IN THE TEES BASIN






FIG. 49

SAMPLE LOCATIONS FOR EACH BEACH AREA


1. CRIMDON BEACH




2.) HARTLEPOOL YACHT CLUB BEACH




3. SEATON CAREW BEACH




4.) REDCAR BEACH



5. MARSKE BEACH



6.) SALTBURN BEACH
[NORTH]



SCALE: 1" = 100'

FIG. 50

CRIMDON BEACH
SPATIAL DISTRIBUTION OF SEDIMENTARY
CHARACTERISTICS





FIG. 51


SALTBURN BEACH (N.)
SHOWING SPATIAL DISTRIBUTION OF SEDIMENTARY CHARACTERISTICS



SCALE: 1" = 100'

FIG. 52

HARTLEPOOL YACHT CLUB BEACH
SPATIAL DISTRIBUTION OF SEDIMENTARY
CHARACTERISTICS



SCALE: 1" = 100'

FIG. 53

MARSKE BEACH SPATIAL DISTRIBUTION OF SEDIMENTARY CHARACTERISTICS




FIG. 54.

REDCAR BEACH **SPATIAL DISTRIBUTION OF SEDIMENTARY** **CHARACTERISTICS**




FIG 55

SEATON CAREW BEACH SPATIAL DISTRIBUTION OF SEDIMENTARY CHARACTERISTICS




FIG. 56

PARTICLE SIZE
TYPICAL CURVES FROM THE SIX BEACH
SITES




FIG. 57

THE LOCATION OF DUNE SAMPLES




FIG. 58

HARTLEPOOL TO CRIMDON DUNE BELT **SPATIAL DISTRIBUTION OF SEDIMENTARY** **CHARACTERISTICS**




FIG. 59

SOUTH GARE DUNE BELT
SPATIAL DISTRIBUTION OF SEDIMENTARY
CHARACTERISTICS




FIG. 60

TYPICAL CURVES OF DUNE SAMPLES




FIG. 61

LOCATION OF RIVER SAMPLES





FIG. 62

THE LOCATION OF FLUVIAL SAMPLES IN THE RIVER TEES



SAND BAR IN THORPE THEWLES BECK, ENGLAND.





FIG. 63

THE LOWER TEES
LOCATION OF RIVER SAMPLES


SAMPLE AREA 1.

THORPE THEWLES BECK
(SANDBANK)




SAMPLE AREA 2.

BILLINGHAM BECK




SAMPLE AREA 3.

SKELTON BECK
(SALTBURN)



SAMPLE AREA 4.

R. TEES (NEASHAM)



SAMPLE AREA 5
S. GARE ESTUARY






FIG. 64

THORPE THEWLES BECK
SPATIAL VARIATION OF SEDIMENTARY CHARACTERISTICS


(A) ϕ MEAN




(B) ϕ STANDARD DEVIATION




(C) ϕ SKEWNESS




(D) ϕ KURTOSIS




(E) % ROCK FRAGMENTS



(F) SHELL CONTENT (GRAINS/1000)



(G) COAL CONTENT (GRAINS/1000)



(H) % HEAVY MINERALS




FIG. 65

BILLINGHAM BECK












FIG. 66


SPATIAL VARIATION OF SEDIMENTARY CHARACTERISTICS

(A) ϕ MEAN




(B) ϕ STD. DEVIATION


SKELTON BECK
(SALTBURN)




(C) ϕ SKEWNESS




(D) ϕ KURTOSIS




(E) % ROCK FRAGMENTS



(F) SHELL FRAGMENTS (GRAINS/1000)



(G) COAL CONTENT (GRAINS/1000)



(H) % HEAVY MINERALS






FIG. 67

**RIVER TEES
(NEASHAM)**
AREAL VARIATION OF SEDIMENTARY CHARACTERISTICS


(A) ϕ MEAN




(B) ϕ DEVIATION (STD.)




(C) ϕ SKEWNESS




(D) ϕ KURTOSIS




(E) % ROCK FRAGMENTS



(F) SHELL FRAGMENTS (GRAINS/1000)



(G) COAL CONTENT (GRAINS/1000)



(H) % HEAVY MINERALS






FIG. 68

SOUTH GARE ESTUARY
AREAL VARIATION OF SEDIMENTARY CHARACTERISTICS

(A) Ø MEAN




(B) Ø STD. DEVIATION



(C) Ø SKEWNESS




(D) Ø KURTOSIS



(E) % ROCK FRAGMENTS



(F) SHELL CONTENT (GRAINS/1000)



(G) COAL CONTENT (GRAINS/1000)



(H) % HEAVY MATERIALS



FIG. 69

TYPICAL CURVES OF RIVER SAMPLES




FIG. 70

THE LOCATION OF OFFSHORE SAMPLES
IN THE TEES BAY




FIG. 71.

OFFSHORE SPATIAL VARIATION OF SEDIMENTARY CHARACTERISTICS




FIG. 72.

OFFSHORE
SPATIAL VARIATION OF SEDIMENTARY CHARACTERISTICS




FIG. 7.3

TYPICAL CURVES OF OFFSHORE SAMPLES




FIG. 74

DROMAN (SUTHERLAND, SCOTLAND)
LOCATION OF SAMPLES ON MARINE AND LAKE SHORES





FIG. 75

TYPICAL CURVES OF SUTHERLAND MARINE AND LACUSTRINE SHORE DEPOSITS PLUS
MARINE DEPOSIT LESS SHELL CONTENT.




FIG.76

**PARTICLE SIZE GRAPH DEMONSTRATING THE EFFECT OF SOURCE MATERIAL
ON BEACH SEDIMENTS**




FIG. 77

ϕ UNITS

BEACH SAMPLES FROM AREAS OUTSIDE THE TEES RIVER BASIN.




FIG. 78

DUNE SAMPLES FROM AREAS OUTSIDE THE TEES RIVER BASIN




FIG.79

RIVER SAMPLES FROM OUTSIDE THE TEES RIVER BASIN .




FIG.80

OFFSHORE SAMPLES FROM AREAS OUTSIDE THE RIVER TEES BASIN




FIG 81

SAND FRACTION FROM BOULDER CLAYS IN NORTHUMBERLAND AND DURHAM.




FIG. 82

FLUVIO - GLACIAL SAMPLES FROM SOUTHEAST ICELAND.




FIG. 83

TYPICAL CURVES OF THE FOUR PRESENT DAY SANDY ENVIRONMENTS
IN THE TEES BASIN




FIG. 84

TYPICAL CURVES OF VARIOUS SANDS AFTER REMOVAL OF SHELL
COAL AND HEAVY MINERALS









FIG. 85

FACTOR LOADING RATIOS OF KNOWN AND UNKNOWN ENVIRONMENTS




 LAKESHORE
 LAKE NEAR SHORE


LACUSTRINE ENVIRONMENT



 OFFSHORE 78% SUCCESSFUL
 DUNE 86% "
 RIVER 100% "
 BEACH 100% "
ENVIRONMENTAL ZONES




DISTRIBUTION OF SAMPLES OF UNKNOWN ORIGIN.



● DUNE
 ○ BEACH
 × RIVER
 ▲ OFFSHORE

DIFFERENTIATION OF ENVIRONMENTS USING FACTOR RATIOS BASED ON PARTICLE SIZE

FACTOR LOADINGS OF KNOWN SAMPLES (BASED ON ALL SEDIMENTARY DATA)



- BEACH
- ▲ DUNE
- RIVER
- OFFSHORE

FIG. 87

PERCENTAGE SILT CONTENT OF THE TEES LAMINATED CLAY DEPOSITS.







FIG. 88

X-RAY DIFFRACTOMETER PATTERNS OF LOBATE, SPHEROIDAL-
ELLIPSOID AND IRREGULAR-PAPILLIFORM GLAUCONITE GRAINS.



AFTER PRATT 1962.

X-RAY DIFFRACTION PATTERNS OF BIOTITE AND GLAUCONITE



AFTER PRATT 1962.

FIG. 90

COMPARISON OF GLAUCONITE CONTAINING DEPOSIT
WITH THE TEES LAMINATED CLAY.






FIG. 91

LOCATION OF BOREHOLES RECORDING LAMINATED CLAY AT DEPTHS
BELOW ORDNANCE DATUM.




• 2' INDICATES DEPTH OF LAMINATED CLAY SURFACE

SUPPOSED MARINE DEPOSIT RECORDED AT SALTBURN BEACH.



SECTION A - B



SECTION C - D




FIG. 93

PARTICLE SIZE CURVES OF THE SUPPOSED MARINE SAND AT SALTBURN





FIG. 94

BRIERTON SAND DEPOSIT, WEST HARTLEPOOL (477299)

LOCATION OF SAMPLES




CLAY DEPOSIT


FIG.95

**GRAPH SHOWING PARTICLE SIZE CURVES OF SAMPLES
FROM BRIERTON**

ARITHMETIC PROBABILITY




**φ SCALE
OFFSHORE GROUP**




**φ SCALE
FLUVIAL GROUP**

ARITHMETIC PROBABILITY



**φ SCALE
BOULDER CLAY GROUP**



NIL

φ SCALE

PARTICLE SIZE CURVES OF SAMPLES FROM ESTON CEMETERY & PARKEND FARM




FIG. 97

PARTICLE SIZE CURVES OF SAND SAMPLES FROM THE ORMESBY BECK VALLEY




FIG. 9/8

**FOUR PARTICLE SIZE CURVES FROM NEWTON BEWLEY SAND
SAMPLES.**









FIG. 99


FACTOR LOADINGS BASED ON PARTICLE SIZE ANALYSIS.




- HURWORTH H. SAMPLES
- HURWORTH
- WYNYARD
- ⊙ WYNYARD BASAL SAND



- COATHAM STOB
- ▲ SANDY FLAT
- SPITTAL FLAT
- CAT FLAT



NEWTON BEWLEY SAND & GRAVEL
PIT



WEST HARTLEPOOL
(BRIERTON - NZ 477299)




FIG. 101

PARTICLE SIZE CURVES OF SAMPLES TAKEN FROM SANDY FLAT (NZ 493158)




FIG. 102

PARTICLE SIZE CURVES OF SAND SAMPLES FROM LEVELLED TERRACES IN
THE TEES BASIN.




FIG.103

LOCATION OF SAMPLE POINTS FOR THE COLLECTION
OF SAND SAMPLES FROM COATHAM STOB.




FIG. 104.

TYPICAL PARTICLE SIZE CURVES OF SAMPLES FROM COATHAM STOB




FIG.105

TYPICAL PARTICLE SIZE CURVES OF SAMPLES FROM HURWORTH

BRICK PIT.




FIG. 106

**TYPICAL PARTICLE SIZE CURVES OF SAMPLES FROM WYNYARD SAND PIT
(NZ 415246)**




FIG. 107



BIBLIOGRAPHY

This bibliography does not attempt to include all works relevant to the subject under discussion but only those which, in the author's opinion, have led to a greater understanding of the relevant topics. Articles are listed under the relevant chapter with a sub-division between those works referenced in the text and other significant works.

Chapter 1

Works Referred to in the Thesis Text

- Agar, R. 1954 The Glacial and Post Glacial Geology of
Middlesborough and the Tees Estuary.
Proc. Yorks. Geol. Soc., V. 29, Pt. 3, pp. 237-253.
- Alexander, C. S. 1953 The Marine and Stream Terraces of the
Capitola Watsonville Area.
University of California pub. in Geog. 10(1),
pp. 1-44.
- Anderson, B. G. 1960 Sorlandet. I Sen-og Post Glacial Tid.
Norg Geol. Unders No. 210, 142 p., Oslo.
- Andrews, J. T. 1966 Pattern of Coastal Uplift and Deglaciation
West Baffin Island, N. W. T.
Geog. Bulletin, V. 8, No. 2, pp. 174-193.
- Andrews, J. T. 1967 Problems in the Analysis of the Vertical
Displacement of Shorelines.
Geog. Bulletin. V. 9, No. 1, pp. 71-74.
- Andrews, J. T. 1968 a Isolines on Maximum Potential Elevations
of the Post Glacial Marine Limit and Post Glacial
Uplift.
Atlas of Canada. Department of Energy, Mines,
and Resources.
- Andrews, J. T. 1968 b Post Glacial Rebound: Similarity and
Prediction of Uplift Curves.
Canadian Journal of Earth Science, pp. 39-47.

- Andrews, J. T. 1969 a The Importance of the Radiocarbon Standard Deviation in Determining Relative Sea Levels and Glacial Chronology From East Baffin Island.
Arctic, Vol. 22, No. 1, 1969, pp. 13-24.
- Andrews, J. T. 1969 b The Shoreline Relations Diagram, Physical Basis and Use in Predicting the Age of Relative Sea Levels; Evidence From Arctic Canada.
Journal of Arctic and Alpine Research, V. 1, pp. 67-78.
- Benema, J. 1954 Holocene Movements of Land and Sea Level in the Coastal Area of the Netherlands.
Geologie en Mijnbouw, N. S., 16(6), pp. 254-262.
- Benema, J., Geve, E. C. W. A., Smits, H., and Wiggers, A. J. 1954
Soil Compaction in Relation to Quaternary Movements of Sea Level and Subsidence of the Land, Especially in the Netherlands.
Geol. en Mijn, 16(6), pp. 173-184.
- Blackburn, K. B. 1952 The Dating of a Deposit Containing an Elk Skeleton Found at Neasham Near Darlington, Co. Durham.
New Phytologist, Vo. 51, pp. 364-377.
- Bloom, A. L. 1960 Late Pleistocene Changes of Sea Level in Southwestern Maine.
Maine Geological Survey.
- Bloom, A. L. 1963 Late Pleistocene Fluctuations of Sea Level and Post Glacial Crustal Rebound in Coastal Maine.
American Journal of Science, V. 261, pp. 862-879.
- Broeker, W. S. 1966 Glacial Rebound and the Deformation of the Shorelines of Proglacial Lakes.
Journal of Geophys. Research, V. 71, pp. 4777-4783.
- Charlesworth, J. K. 1957 The Quaternary Era.
2 Vols., E. Arnold, Edinburgh.
- Churchill, D. B. 1965 The Displacement of Deposits Formed at Sea Level 6,500 Years Ago in Southern Britain.
Quaternaria, Vol. 7, pp. 239-250.

- Cullingford, R. A., Smith, D. E. 1966 Late Glacial Shorelines in Eastern Fife.
Inst. Brit. Geog., Spec. No. on Vertical Displacement of Shorelines, No. 39, pp. 31-51.
- Curray, J. 1961 Late Quaternary Sea Level; a Discussion.
Bulletin American Geol. Soc., V. 72, pp. 1707-1712.
- Daly, R. A. 1920 A Recent World Wide Sinking of Ocean Level.
Geol. Mag., 57, pp. 246-261.
- Daly, R. A. 1934 The Changing World of the Ice Age.
Yale University Press, New Haven, 271 pp.
- Donn, W. L., Farrand, W. R., Ewing, M. E. 1962 Pleistocene Ice Volumes and Sea Level Lowering.
Journal of Geology, V. 70, pp. 206-214.
- Donner, J. J. 1959 The Late and Post Glacial Beaches in Scotland.
Ann. Acad. Sci. Fenn., Ser. A, III, 53.
- Donner, J. J. 1963 The Late and Post Glacial Raised Beaches in Scotland II.
Ann. Acad. Sci. Fenn., Ser. A, 68.
- Donner, J. J. 1965 Shoreline Diagrams in Finnish Quaternary Research.
Baltica, Vol. 2, pp. 11-20.
- Fairbridge, R. W. 1958 Dating the Latest Movements of the Quaternary Sea Level.
Trans N. Y. Acad. Sci., Series ii, V. 20, No. 6, pp. 471-482.
- Fairbridge, R. W. 1961 Eustatic Changes in Sea Level.
Physics and Chemistry of the Earth, Vol. 4, pp. 99-185.
- Fairbridge, R. W. 1962 World Sea Level and Climatic Changes.
Quaternaria, Vol. 6, pp. 111-134.
- Farrand, W. G. 1962 Post Glacial Uplift in North America.
Am. Journ. Sci., V. 260, pp. 181-199.

- Farrand, W. G. 1965 The Deglacial Homicycle.
Geol. Rundschau, V. 54, pp. 385-398.
- Fisk, H. N. 1939 Depositional Terrace Slopes in Louisiana.
Journ. Geomorph., V. 2, pp. 181-200.
- Fisk, H. N., McFarlan, E. 1955 Late Quaternary Deposits of the
Mississippi River.
Geol. Soc. Am. Bull., Spec. Paper 62, pp. 279-302.
- Gillberg, G. 1952 Marina Gransen i Vastra Sverige.
Geol. Foren Stockholm Forand, 74, pp. 71-103
(English Abstract).
- Godwin, H. 1943 Coastal Peat Beds of the British Isles and
North Sea.
Journal of Ecology, V. 31, pp. 199-247.
- Godwin, H. 1945 Coastal Peat Beds of the North Sea Region as
Indices of Land and Sea Level Changes.
New Phytologist, V. 44, pp. 29-69.
- Godwin, H., Suggate, R. P., Willis, E. H. 1958 Radiocarbon
Dating of the Eustatic Rise in Sea Level.
Nature, V. 181, p. 1518.
- Godwin, H. 1960 Radiocarbon Dating and Quaternary History in
Britain.
Proc. Royal Soc., London, B, 153, pp. 287-320.
- Green, C. 1961 East Anglian Coast Line Levels Since Roman Times.
Antiquity, 35(137), pp. 21-28.
- Grosvald 1969 Institute of Academy of Sciences, U. S. S. R.
Soviet Research in Franz Josef Land.
- Gutenberg, B. 1941 Changes in Sea Level, Post Glacial Uplift
and Mobility of the Earth's Interior.
Bull. Geol. Soc. Am., Vol. 52, pp. 721-772.
- Hogbom, A. G. 1919 Eine graphische Darstellung der Spatquartaren
Niveau Verand er Ungen Fennoskandias.
Bull. Geol. Instit. of Upsala, XVI.

- Jelgersma, S. 1961 Holocene Sea Level Changes in the Netherlands. Medelingen Geologische Stichting, Series C, Vol. VI, No. 7, Maastricht.
- Jelgersma, S. 1966 Sea Level Changes During the Last 10,000 Years, IN International Symposium on World Climate From 8000 to 0 B.C. London Royal Meteorol. Soc., pp. 54-71.
- Johnson, W. A. 1946 Glacial Lake Agassiz; with Special Reference to the Mode of Deformation of the Beaches. Geol. Survey Canada Bull. No. 7, 20 pp.
- Kaizuka, S. 1959 The Physiographic Development of the Kanto Plain Tephrochronological Study. Proc. Inter Geog. Union, Reg. Conf. in Japan.
- Kaariainen, E. 1953 On the Recent Uplift of the Earth's Crust in Finland. Fennia, V. 77, No. 2, 106 pp.
- Kendall, P. F. 1902 A System of Glacier Lakes in the Cleveland Hills. Quat. Journ. Geol. Soc., pp. 471-571.
- King, C. A. M. 1966 Techniques in Geomorphology. Arnold, Edinburgh.
- Kirk, W., Rice, R. J., Synge, F. M. 1966 Deglaciation and Vertical Displacement of Shorelines in Wester and Easter Ross. Institute of Brit. Geog., Spec. No. on Vertical Displacement of Shorelines, No. 39, pp. 65-78.
- Lawson, A. C. 1940 Isostatic Control of Fluctuations of Sea Level. Science, Vol. 92, pp. 162-164.
- LeBlanc, R. J., Bernard, H. A. 1954 Resume of Recent Geological History of the Gulf Coast. Geol. en Mijnbouw, N. S., 16, No. 6, pp. 185-194.
- Løken, O. 1965 Post Glacial Emergence at the South End of Imigsuin Fiord, Baffin Island, N. W. T. Geog. Bull., V. 7, pp. 243-258.

- McCann, S. B. 1961 The Raised Beaches of Western Scotland.
Unpublished Ph.D. dissertation, University of
Cambridge.
- McCann, S. B. 1964 The Raised Beaches of North-East Islay and
Western Jura, Argyll.
Trans. Inst. Brit. Geog., V. 35, pp. 1-16.
- McCann, S. B. 1966 The Main Post Glacial Raised Shoreline of
Western Scotland from the Firth of Lorne to
Loch Broom.
Trans. Inst. Brit. Geog., Spec. No. 39, Vertical
Displacement of Shorelines, pp. 87-99.
- Movius, H. J., jnr. 1953 Graphic Representation of Post Glacial
Changes of Level in North-East Ireland.
Amer. Journ. Sci., V. 251, No. 10, pp. 697-740.
- Nanson, F. 1927 The Earth's Crust, Its Surface Forms and
Isostatic Adjustment.
Avhandlingar Det Norske-Videnskaps Akedemi
(Matematisk-Natur Viden Skapelig Klasse, Oslo),
117 pp.
- Nye, J. F. A. 1959 A Method of Calculating the Thickness of
Ice Sheets.
Nature, V. 169, pp. 529-530.
- Pannekoek, A. J. 1954 Tertiary and Quaternary Subsidence in
the Netherlands.
Geol. en Mij., N. S., 16, pp. 156-164.
- Sauramo, M. 1941 Fennoskandien in Quatar.
Reg. Geol. der Erde, Vol. 1, 22 pp.
- Sauramo, M. 1955 Land Uplift with Hinge Lines in Fennoscandia.
Ann. Acad. Sci. Fenn., Ser. A, III, No. 44.
- Sauramo, M. 1958 Die Geschichte der Ostsee.
Ann. Acad. Sci. Fenn., Ser. A, III, No. 51.
- Raistrick, A. 1931 The Glaciation of Northumberland and Durham.
Proc. Geol. Assoc., 42, pp. 281-291.
- Reid, C. 1913 Submerged Forests.
Cambridge University Press.

- Schofield, J. C. 1960 Sea Level Fluctuations During the Last 4,000 Years as Recorded by a Chenier Plain, Firth of Thames, New Zealand. N. Z. Journ. of Geol. and Geophys., V. 3, pp. 467-485.
- Schofield, J. C. 1964 Post Glacial Sea Levels and Isostatic Uplift. N. Z. Journ. Geol. and Geophys., V. 7, pp. 359-370.
- Schofield, J. C. 1963 Symposium on Changes of Climate in the Post Glacial Period. Proc. N. Z. Ecolog. Soc., V. 10, pp. 21-52.
- Shepard, F. P. 1963 Thirty-Five Thousand Years of Sea Level, IN Essays in Marine Geology in Honour of K. O. Emery. Los Angeles, University of Southern California.
- Shepard, F. P., Suess, H. E. 1956 Rate of Post Glacial Rise in Sea Level. Science, V. 123, pp. 1082-1083.
- Shotton, F. W. 1967 The Problems and Contributions of Methods of Absolute Dating Within the Pleistocene Period. Quarterly Journal of the Geol. Soc., Vol. 122, pp. 357-383.
- Sissons, J. B. 1962 A Re-Interpretation of the Literature on Late Glacial Shorelines in Scotland with Particular Reference to the Forth Area. Trans Edin. Geol. Soc., Vol. 19, pp. 83-99.
- Sissons, J. B. 1963 Scottish Raised Shoreline Heights with Particular Reference to the Forth Valley. Geografiska Annaler, Vol. 45, pp. 180-185.
- Sissons, J. B., Smith, D. E. 1965 Raised Shorelines Associated with the Perth Readvance in the Forth Valley and Their Relation to Glacial Isostasy. Trans. Royal Soc. Edin., Vol. 46, pp. 143-168.
- Sissons, J. B., Smith, D. E., Cullingford, R. A. 1966 Late Glacial and Post Glacial Shorelines in South-East Scotland. Trans. Inst. Brit. Geog., Spec. No. Vertical Displacement of Shorelines, pp. 9-18.

- Sissons, J. B. 1966 Relative Sea Level Changes Between 10300 and 8300 B.P. in Part of the Carse of Stirling. Trans. Inst. Brit. Geog., Spec. No. 39, Vertical Displacement of Shorelines, pp. 19-29.
- Smith, D. E. 1965 Late and Post Glacial Changes of Shoreline on the North Side of the Forth Valley and Estuary. Unpublished Ph.D. Thesis, University of Edinburgh.
- Smith, D. E. 1968 Post Glacial Displaced Shorelines in the Surface of the Carse Clay on the North Bank of the River Forth in Scotland. Zeitschifte für Geomorphologie, Vol. 12, pp. 388-408.
- Stephens, N. 1963 Late Glacial Sea Levels in N. E. Ireland. Irish Geography, Vol. 4, pp. 345-359.
- Stephens, N. 1966 Late Glacial and Post Glacial Shorelines in Ireland and South-West Scotland. Spec. Geol. Soc. Am. Papers, Int. Stud. Quaternary VII, Congress Int. Ass. Quaternary Research, Boulder, Colorado.
- Stephens, N., Synge, F. M. 1965 Late Pleistocene Shorelines and Drift Limits in North Donegal. Proc. R. Ir. Acad., 64B, pp. 131-153.
- Stephens, N., Synge, F. M. 1966 Pleistocene shorelines. In G. H. Dury, ed. Geomorphological Essays, pp. 1-51.
- Synge, F. M. 1966 The Relationship of the Raised Strand Lines and Main End-Moraines on the Isle of Mull and in the District of Lorn, Scotland. Proc. Geol. Assoc., Vol.
- Synge, F. M., Stephens, N. 1966 Late and Post Glacial Shorelines and Ice Limits in Argyll and N. E. Ulster. Trans. Inst. Brit. Geog., Spec. No. 39, Vertical Displacement of Shorelines, pp. 101-125.
- Umbgrove, J. H. F. 1950 A Country Below Sea Level. Symphony of the Earth, Martinus Nijhoff, The Hague, pp. 31-54.

- Van Straaten, L. M. J. U. 1954 Radiocarbon Datings and Changes of Sea Level at Velzen (Netherlands).
Geol. en Mijnb, N. S. 16, pp. 247-254.
- Van Veen, J. 1954 Tide Gauges, Subsidence Gauges and Flood Stones in the Netherlands.
Geol. en Mijnb, N. S. 16, pp. 214-222.
- Vening-Meinesz, F. A. 1954 Crustal Warping in the Netherlands.
Geol. en Mijnb, N. S. 16, p. 207.
- Weertman, J. 1961 Equilibrium Profile of Ice Caps.
Journ. Glaciol., V. 3, pp. 953-964.
- Weertman, J. 1966 Effect of Basal Water Layer on Dimensions of Ice Sheets.
Journ. of Glaciol., V. 6, pp. 191-208.
- West, R. G. 1963 Problems of the British Quaternary.
Proc. Geol. Assoc., Vol. 74, Pt. 2, pp. 147-186.
- West, R. G. 1968 Pleistocene Geology and Biology.
Longman's, London.
- Wiggers, A. J. 1954 Compaction of Sediments.
Geol. en Mijnb., N. S. 16, pp. 179-184.
- Wright, W. B. 1914 The Quaternary Ice Age.
Macmillan, London.

Other Significant Works

- Airy, G. B. 1855 On the Computation of the Effect of the Attraction of Mountain Masses as Disturbing the Apparent Astronomical Latitude of Stations in Geodetic Surveys.
Proc. Royal Soc., V. 7, pp. 240-242.
- Antews, E. 1939 Late Quaternary Upwarps of Northeastern North America.
Journ. of Geol., Vol. XLVII, pp. 707-720.

- Baden-Powell, D. F. W. 1963 Isostatic Recovery in Scotland.
Nature, Vol. 199, pp. 546-547.
- Barrell, J. 1915 Factors in Movements of the Strand Line and
Their Results in the Pleistocene and Post
Pleistocene.
Amer. Journ. Sci., V. 40, pp. 1-22.
- Blanc, A. C. 1937 Low Levels of the Mediterranean Sea During
the Pleistocene Glaciation.
Quat. Journ. Geol. Soc., Vol. 93, pp. 621-651.
- Bott, M. H. P. 1962 A Simple Criterion For Interpreting
Negative Gravity Anomalies.
Geophysics, Vol. 27, pp. 376-381.
- Bott, M. H. P. 1964 a The Deep Sea Structure of the Northern
Irish Sea,
IN Submarine Geology and Geophysics.
Colston Research Papers, pp. 179-204.
- Bott, M. H. P. 1964 b Formation of Sedimentary Basins by
Ductile Flow of Isostatic Origin in the Upper
Mantle.
Nature, Vol. 201, pp. 1082-1084.
- Bowie, W. 1912 Effect of Topography and Isostatic Compensation
Upon the Intensity of Gravity.
U. S. Coast and Geodetic Survey, Spec. Pub. No. 12,
Geodesy, Washington, D. C.
- Bowie, W. 1921 The Relation of Isostasy to Uplift and Subsidence.
Am. Journ. of Sci. II.
- Broeker, W. S. 1962 The Contribution of Pressure Induced Phase
Changes to Glacial Rebound.
Journ. Geophys. Research, V. 67, pp. 4837-4842.
- Brogger, W. C. 1901 Om de Senglaciale og Postglaciale Nivator
and Ringar i Kristiania Felgt (On the Late and
Post Glacial Changes of Level in the Kristiana
Region) with English Summary.
Norg Geol. Unders, V. 31, 731 pp.
- Chamberlin, R. T. Isostasy from the Geological Point of View.
Journ. of Geol., V. 39, pp. 1-23.

- Croll, J. 1866 On the Physical Cause of the Submergence and Emergence of the Land During the Glacial Epoch. London-Edinburgh and Dublin Phil. Mag. and Journal of Science, 4th Series, V. 31, pp. 301-306.
- DeGeer, G. 1892 Quaternary Changes of Level in Scandinavia. Bull. Am. Geol. Soc., V. 3, pp. 65-68.
- Farrand, W. R., Gajda, R. T. 1961 Isobases of Maximum Post Glacial Submergence in Glaciated North America, IN I. N. Q. U. A., 6th Congress Abstract of Papers, p. 178.
- Godwin, H., Baden-Powell, D. F. W. 1963 Isostatic Recovery in Scotland. Nature, V. 199, pp. 1277-1278.
- Griggs, D. T. 1960 Isostatic Compensation and Origin of Continents and Mountains. Trans. Amer. Geophys. Union, V. 41, pp. 166-167.
- Gunn, R. 1949 Isostasy Extended. Journ. of Geol., V. 57, pp. 263-279.
- Hallam, A. 1963 Major Epeirogenic and Eustatic Changes Since the Cretaceous and Their Possible Relationship to Crustal Structure. Am. Journ. Sci., V. 261, pp. 397-423.
- Harrison, W., Lyon, C. J. 1963 Sea Level and Crustal Movements Along the New England Acadian Shore, 4500 - 3000 B.P. Journ. Geol., Vol. 71, pp. 96-108.
- Haug, E. 1900 Les Geosynclinause et les Aires Continentales. Bull. Geol. Soc. Amer., V. 28, pp. 617-711.
- Hedberg, H. D. 1936 Gravitational Compaction of Clays and Shales. Am. Journ. of Sci., V. 31, pp. 241-285.
- Heiskanen, W. A., Vening-Meinesz, F. A. 1958 The Earth and Its Gravity Field. McGraw-Hill Book Co., Inc.
- Hicks, S. D., Shofnos, W. 1965 The Determination of Land Emergence from Sea Level Observations in South-East Alaska. Journ. Geophys. Research, V. 70, pp. 3315-3320.

- Hutton, J. 1785 The Theory of the Earth.
Trans. Royal Soc. Edin., Pamphlet.
- Ives, J. 1967 Review of the I. B. G. Special No. 39, Vertical
Displacement of Shorelines.
Geog. Bulletin, Vol. 9, No. 1, pp. 76-77.
- Jamieson, T. F. 1882 On the Causes of the Depression and
Re-Elevation of the Land During the Glacial
Period.
Geol. Mag., V. 9, pp. 400-407.
- Jamieson, T. F. 1887 On Some Changes of Level During the Glacial
Period and Their Supposed Cause.
Geol. Mag.
- Jamieson, T. F. 1905 Some Changes of Level in the Glacial
Period.
Geol. Mag., Ser. 5, No. 2, pp. 484-490.
- Jamieson, T. F. 1908 On Changes of Level and the Production of
Raised Beaches.
Geol. Mag., N. S., Vol. 5, pp. 206-209.
- Jones, O. T. 1934 Some Observations on Recent Geological
Movements of the British Coastline.
Antiquity, V. 8, p. 303.
- Kaye, C. A., Barghoorn, E. S. 1964 Late Quaternary Sea Level
Change and Crustal Rise at Boston, Massachusetts,
with Notes on Auto Compaction of Peat.
Bull. Geol. Soc. Am., V. 75, pp. 63-80.
- Kendall, P. F. 1923 On Quaternary Isostatic Readjustments in
N. E. Europe.
Rep. Brit. Assoc., p. 441.
- Kuenen, Ph. H. 1954 Eustatic Changes of Sea Level.
Geol. en Mijnb., N. S. 16, No. 6, pp. 148-155.
- Lyustikh, E. N. 1957 Isostasy and Isostatic Hypotheses.
Soviet Research in Geophysics, in English
Translation, Vol. 2, (Moscow), American Geophys.
Union.
- Maclaren, C. 1842 The Glacial Theory of Prof. Agassiz.
Am. Journ. Sci., V. 42, pp. 346-365.

- Penck, A. 1934 Theorie der Bewegung der Strand Linie.
Sitzber Preuss, Akad. Wiss. Berlin, Phys. Math.,
K. 1, p. 19.
- Pratt, J. H. 1855 On the Attraction of the Himalaya Mountains
and of the Elevated Region Beyond Them, Upon the
Plumb Line in India.
Proc. Royal Soc., V. 7, pp. 176-182.
- Ramsay, W. 1923 On Relations Between Crustal Movements and
Variations of Sea Level During the Late Quaternary
Time Especially in Fennoscandia.
Comm. Geol. Fin. Bull., Vol. 60-69, pp. 1-39.
- Ramsay, W. 1924 Crustal Movements and Variations of Sea Level.
Bull. Comm. Geol. Fin., No. 66, 39 pp.
- Roden, G. I. 1963 Sea Level Variations at Panama.
Journ. Geophys. Research, V. 68, pp. 5701-5710.
- Roden, G. I. 1966 Low Frequency Sea Level Oscillations Along
the Pacific Coast of North America.
Journ. of Geophys. Research, V. 71, No. 20,
pp. 4755-4776.
- Suess, E. 1888 Das Antlitz der Erde.
Vol. II, Vienna.
- Taylor, A. 1853 On Changes of the Sea Level Effected by Existing
Physical Causes During Stated Periods of Time.
Abstract in Quat. Journ. Geol Soc., V. 9, pp. 47-49.
- Valentin, H. 1953 Present Vertical Movements in the British
Isles.
Geog. Journal, Vol. 119, pp. 299-305.
- Von Post, L. 1948 New Aspects on the Late Quaternary Displacement
of the Swedish Sea Shore.
Internat. Geol. Cong., 18th Session, Great Britain,
pp. 219-225.
- Wright, W. B. 1925 Three Short Papers on Isostasy.
Geol. Mag., V. 62, pp. 227-234.
- Wright, W. B. 1934 The Isokinetic Theory of Glacially Controlled
Shorelines.
Comptes Rendus du Congres Internationale de
Geographie Varsovie, Sect. II, pp. 534-553.

CHAPTER 2

BIBLIOGRAPHY

Publications discussed in the text are listed below. The reader is referred to the following sources, from which the author made his survey of additional works:

- (1) Annotated Bibliography of Quaternary Shorelines, 1945-1964.
Richards, H. G., and Fairbridge, R. W., INQUA, 1965.
- (2) Geo. Abstracts, Volume A, University of East Anglia.
- (3) Smith, D. E. (Unpublished Ph.D., University of Edinburgh),
(Early History), 1965.

(Publications discussed and included in the bibliography for Chapter 1 are denoted "op. cit.").

- Addicott, W. O. 1963 Interpretation of the Invertebrate Fauna from the Upper Pleistocene Battery Formation near Crescent City, California.
Proc. Calif. Acad. Sci., V. 31, No. 13, pp. 341-347.
- Addicott, W. O. 1964 A Late Pleistocene Invertebrate Fauna from South-western Oregon.
Journal of Paleontology, V. 38, Pt. 4, pp. 650-661.
- Anderson, F. W. 1947 The Fauna of the '100 Foot Beach' Clays.
Trans. Edin. Geol. Soc., Vol. 14, pp. 220-229.
- Arnold, B. A. 1957 Late Pleistocene and Recent Changes in Land Forms, Climate, and Archaeology in Central Baja, California.
Univ. Calif. Pub. in Geogr., 10(4), pp. 201-318.
- Auer, V. 1950 Las capas Volcanicas Como Base de la Cronologia Postglacial de Fuego-Patagonia.
Ministerio de Agricultway Ganaderia, Pub. 9, Buenos Aires.

- Auer, V. 1959 The Pleistocene of Fuego-Patagonia, Part III, Shoreline Displacements.
Ann Acad. Sci. Fenn, Ser. A, No. 60, Helsinki.
- Avnimelech, M. 1950 On the Finding of Elephas Bones in a Boring near Yavney, Israel.
Eclogae Geologicae Helvetiae, 43(2).
- Baldwin, E. M. 1959 Geology of Oregon.
University of Oregon, 136 pp., Eugene, Oregon.
- Bandy, O. L. 1953 Ecology and Paleoecology of Some California Foraminifera.
Journal of Paleontology, 27(2), pp. 200-203.
- Barrett, W. 1962 Emerged and Submerged Shorelines of the Dominican Republic.
Revista Geographica, 30(1), pp. 51-77.
- Bergsten, F. 1930 Changes of Sea Level on the Coasts of Sweden.
Geog. Annaler, V. 12, pp. 21-55.
- Birkenmajer, K. 1958 Preliminary Report on the Raised Marine Features in Hornsund Vestspitzbergen.
Bull. Acad. Polonaise des Sci. Ser. Sci. Chim. Geol. et Geogr., 6(2), pp. 151-157.
- Bjorsjo, N. 1953 Nytt Sub fossil Fynd Av Val Pa Vastkusten.
(A new Find of Sub-fossil Whale on the West Coast of Sweden.)
Geol. Foren. i Stockholm, For. V. 75, pp. 510-513.
- Blanc, A. C., Segre, A. G., Tongiorgi, E. 1953 Le Quaternaire de l'Agro Pontino.
IV Congr., INQYA, Rome.
- Boillot, G. 1964 Geologie de la Manche Occidentale.
Ann de l'Institut Oceanographique, 42(1), pp. 1-320.
- Bradley, W. H. 1953 Age of Intertidal Tree Stumps at Robinhood, Maine.
Amer. Journ. Sci., V. 251, pp. 543-546.
- Brady, G. S. 1873 On the Method of Collecting and Preserving Entomostraca and Other Microzoa.
Nature, Vol. 8.

- Brodie, J. W. 1955 Sedimentation in Lyttelton Harbour, South Island, New Zealand.
N. Z. Jour. Sci. Tech., 363(6), pp. 603-621.
- Brown, E. H. 1952 The 600 Foot Platform in Wales.
Proc. 17th Inter Geog. Cong., pp. 304-312, Washington.
- Burnaby, T. P. 1950 The Tubular Chalk Stacks of Sheringham.
Proc. Geol. Assoc. Lond, 61(4), pp. 226-241.
- Cadilla, J. F. 1960 Geology of the Carolina Clay Deposit in Puerto Rico.
Trans. 2nd Caribbean Geol. Conf., pp. 147-149.
- Caldenius, C., and Linnman, G. 1949 A Late Quaternary Regression and Transgression at Halmstad, Southern Sweden.
Sveriges Geologiska Unders, Ser. C., p. 502.
- Cayeux, L. 1941 Causes Anciennes et Causes Actuelles en Geologie.
Masson, Paris, 81 pp.
- Chappell, J. 1967 Recognizing Fossil Strand Lines from Grain Size Analyses.
Journ. Sed. Pet., Vol. 37(1), pp. 157-165.
- Churchill, D. M. 1959 Late Quaternary Eustatic Changes in the Swan River District.
Journ. Royal Soc. W. Aust. (Perth), V. 42, pp. 53-55.
- Clements, T. and Dana, S. W. 1944 Geologic Significance of a Coarse Marine Sediment from near Santa Cataline Isle, California.
Journ. Geol., 52, pp. 351-354.
- Claridge, G. G. C. 1961 Mineralogy and Origin of the Yellow-Brown Sands and Related Soils.
N. Z. Jour. Geol. and Geophyz, 4(1), pp. 48-72.
- Cloud, P. E., Schmidt, R. G., Burke, H. W. 1959 Geology of Saipan Mariana Islands, Pt. 4, Submarine Topography and Shoalwater Ecology.
U. S. Geol. Survey, Prof. Paper, 280-K.
- Coats, R. R. 1956 Geology of Northern Adak Island, Alaska.
U. S. Geol. Surv. Bull, 1028-C.

- Coats, R. R., Nelson, W. H., Lewis, R. Q., Powers, H. A. 1961
Geologic Reconnaissance of Kiska Island, Aleutian
Islands, Alaska.
U. S. Geol. Surv. Bull., 1028-R.
- Colquhoun, D. J. 1962 On Surficial Sediments in Central South
Carolina: A Progress Report.
Geol. Notes, Colombia, S. C., 6(6), pp. 63-80.
- Colquhoun, D. J. 1962 Wicomico Shoreline in Orangeburg, Dorchester,
and Berkeley Counties.
Geol. Notes, Colombia, S. C., 5(4), pp. 43-50.
- Cooper, L. H. N. 1948 A Submerged Ancient Cliff near Plymouth.
Nature, Vol. 161, p. 280.
- Cooke, C. W. 1945 Geology of Florida.
Fla. Geol. Surv. Bull. 29.
- Cooke, C. W. 1951 Geologic Map of Prince Georges County (Maryland)
and the District of Columbia.
Md. Department Geol., Mines and Water Resources.
- Cooke, C. W. 1952 Sedimentary Deposits of Prince Georges County
(Maryland) and the District of Columbia.
Md. Dept. Geol. Mines and Water Resources Bull. 10,
pp. 1-53.
- Cooke, C. W. 1958 Pleistocene Shorelines in Maryland.
Bull. Geol. Soc. Am., 69(9), pp. 1187-1190.
- Crawford, A. R. 1963 Quaternary Sedimentary Breccias and
Emerged Offshore Bars near Point Lowly.
Quat. Geol. Notes, Geol. Surv., S. A. Adelaide,
Vol. 5, pp. 1-2.
- Baczyk, J. 1963 Genese de la Presquile de Hel sur la Base du
Developpement du Golf de Gdansk.
Baltica, Vol. 1, pp. 174-188.
- Dannstedt, G. 1948 Bottentopografien i Sodra Kalmarsund.
Geog. Annaler, V. 29, pp. 1-19.
- Davies, G. L. 1958 Irish Erosion Surfaces--A Statistical Analysis.
Adv. of Sci., Vol. 14, pp. 385-388.

- Davies, G. L. 1960 Irish Erosion Surfaces.
Geol. Mag., Vol. 98(2), pp. 172-173.
- Davies, G. L. 1963 Platforms Developed in the Boulder Clay of
the Coastal Margins of Counties Wicklow and Wexford.
Irish Geog., Vol. 4, pp. 107-116.
- Devyatova, E. I. 1960 Marine Interglacial Deposits in the Onega
River Basin.
Akad. Nauk. Cokladi, Vol. 125, pp. 1-6.
- Dibblee, T. W., jnr. 1950 Geology of South-western Santa Barbara
County, California.
Calif. Div. of Mines, Bull. 150.
- Doeglas, D. J. 1950 Old Beaches in the Mediterranean.
18th Int. Geol. Cong., London, Pt. 8, pp. 16-17.
- Donner, J. J. 1952 On the Early Post glacial Shoreline Displacement
in South-East Finland.
Ann. Acad. Sci. Fenn., Ser. A, 3, p. 29.
- Donner, J. J. 1957 The Post Glacial Shoreline Displacement
in the Kuopi District.
Ibid., p. 49.
- Donner, J. J. 1959 Op. cit.
- Donner, J. J. 1963 Op. cit.
- Donner, J. J. 1965 Op. cit.
- Downing, R. A. 1959 A Note of the Crag in Norfolk.
Geol. Mag., V. 96(1), pp. 81-86.
- Dyvyetov, E. I., and Losyev, E. I. 1964 Stratigraphy and
Paleontology of the Quaternary Period of the Basin of
the Mezen River.
- Earp, J. R., Francis, E. H., Read, W. A. 1962 Discussion of
J. B. Sisson's Paper on Late Glacial Shorelines.
Trans. Edin. Geol. Soc., 19(2), pp. 216-220.
- Emerson, W. K. 1960 Shell Middens of San Jose Island (Baja,
California).
Amer. Mus. Nat. Hist. Novitates, 2013, pp. 1-9.

- Emery, K. O. 1962 Marine Geology of Guam.
U. S. Geol. Survey, Prof. Paper, 403-B.
- Emery, K. O. and Niino, H. 1963 Sediments of the Gulf of
Thailand and Adjacent Continental Shelf.
Bull. Geol. Soc. Amer., 74(5), pp. 541-554.
- Emery, K. O. and Terry, R. D. 1956 A Submarine Slope off
Southern California.
Journ. of Geol., 64(3), pp. 271-280.
- Evers, W. 1962 The Problem of Coastal Genesis with Special
Reference to the "Strandflat," the "Banks," or
"Grounds," and "Deep Channels" of the Norwegian and
Greenland Coasts.
Journal of Geol., V. 70(5), pp. 621-630.
- Ewing, J., Le Pichon, X., Ewing, M. 1963 Upper Stratification
of Hudson Apron Region.
Journ. Geophys. Research, V. 68(23), pp. 6303-6316.
- Fairbridge, R. W., and Stewart, H. B. 1960 Alexa Bank, A
Drowned Atoll on the Melanesian Border Plateau.
Deep Sea Research, Vol. 7, pp. 100-116.
- Federov, P. V. 1943 The Caspian Terraces of the South Mangishlak
Shore.
Doklady Acad. Nauk, U. S. S. R., 9.
- Federov, P. V. 1948 Sea Terraces of the Eastern Shore of the
Caspian Sea.
Doklady Acad. Nauk, U. S. S. R., 59(9).
- Federov, P. V. 1952 Stratigraphic Scheme of the Caspian
Quaternary Deposits.
Moscow Bull., Soc. Exper. in Nature, Geol. Div. 4.
- Federov, P. V., and Leontiev, O. K. 1953 On the History of the
Caspian Sea in the Late and Post Khvalin Time.
Izvestia, Acad. Nauk, U. S. S. R. Geogr., Ser. 4.
- Feyling-Hanssen, R. W. 1964 Foraminifera in Late Quaternary
Deposits from the Oslo Fjordaren.
Norg Geol. Unders, V. 225, pp. 1-383.
- Flinn, D. 1964 Coastal and Submarine Features Around the
Shetland Islands.
Proc. Geol. Assoc. Lond, 75(3), pp. 321-339.

- Folk, R. L. 1967 A Review of Grain Size Parameters.
Sedimentology, V. 6, pp. 73-94.
- Fray, C., and Ewing, M. 1963 Wisconsin Sea Level as Indicated
in Argentina Continental Shelf Sediments.
Proc. Acad. Nat. Sci. Phila., 115, pp. 113-126.
- Friedman, G. M. 1961 Distinction Between Dune, Beach and River
Sand from Textural Characteristics.
Journ. Sed. Pet., V. 31, pp. 514-529.
- Gadd, N. R. 1960 Surficial Geology of the Becancour Map Area,
Quebec.
Geol. Survey Canada, Paper 59, p. 8.
- Gallagher, E. W. 1935 Glauconite Genesis.
Bull. Am. Geol. Soc., V. 46(9), pp. 1351-1366.
- Ganeshin, G. S., and Chemekov, J. F. 1960 Quaternary Stratigraphy
and Paleography in the North-East and Far East of the
U. S. S. R.
IN Chronology and Climatology of the Quaternary.
Akad. Nauk, U. S. S. R., pp. 173-182.
- Geikie, A. 1863 On the Phenomena of the Glacial Drift of Scotland.
Geol. Soc. Glasgow Trans, V. 1(2), 190 pp.
- Gill, D. G. 1960 Summary of the Report of the A. N. Z. A. A. S.
Sections C and P, Committee for the Investigation of
Quaternary Strand Line Changes to the 1959 Perth
Congress.
Aust. Journ. Sci., 23(3), pp. 74-76.
- Gillberg, G. 1952 Marina Gransen i Vastra Sverige.
Geol. Foren Stockholm Forhand, V. 74, pp. 71-103.
- Goreau, T. F. 1961 The Structure of the Jamaican Reef Communities.
Geological Aspects.
Final Progress Report to Biol. Branch, Off. Nav. Res.
N. Y. Zoolog. Soc.
- Gould, H. R., and Stewart, R. H. 1955 Continental Terrace
Sediments in the North-East Gulf of Mexico.
IN "Finding Ancient Shorelines."
Soc. Econ. Pal. and Min., Spec. Pub. 3, pp. 2-19.

- Green, J. F. N. 1946 The Terraces of Boumemouth, Hants.
Proc. Geol. Assoc. Lond., V. 57, pp. 82-101.
- Green, J. F. N. 1949 The History of the River Dart., Devon.
Proc. Geol. Assoc. Lond., V. 60(2), pp. 105-124.
- Gutenberg, B. 1941 Changes in Sea Level, Post Glacial Uplift
and Mobility of the Earth's Interior.
Bull. Am. Geol. Soc., V. 52, pp. 721-772.
- Hack, J. T. 1955 Geology of the Brandy Wine Area and Origin of
the Upland of Southern Maryland.
U. S. Geol. Survey, Prof. Paper, 267-A.
- Harrison, W. 1962 Pleistocene Record in the Sub-surface of the
Norfolk Area, Virginia.
Va. Acad. Sci., Gaudebook, pp. 35-61.
- Hawkins, A. B. 1962 The Buried Channel of the Bristol Avon.
Geol. Mag., 99(4), pp. 369-374.
- Heezen, B. C., Tharp, M., Ewing, M. 1959 The Floors of the
Oceans. The North Atlantic.
Geol. Soc. Am., Spec. Paper 65.
- Heiskanen, W. A., Vening Meinesz, F. A. 1959 The Earth and its
Gravity Field.
McGraw-Hill.
- Hickox, C. F., jnr. 1962 Pleistocene Geology of the Central
Annapolis Valley, Nova Scotia.
Nova Scotia Dept. Mines Mem., V. 5, pp. 1-36.
- Holtedahl, O. 1955 Norges Geologi.
Norg. Geol. Unders, 164.
- Hopgood, A. M. 1961 The Geology of the Cape Rodney-Kawan District,
Auckland.
N. A. Journ. Geol., Geophys., 4(2), pp. 205-230.
- Jahn, A. 1959 Post glacialny rozwój wybrzezy Spitz Bergebu.
Czasopismo Geograficzne, 30, Warsaw.
- Jahns, R. H. 1954 Geology of the Pensinsular Range Province
Southern California and Baja, California.
California Div. Mines Bull. 170(2), pp. 53-63.

- Jamieson, T. F. 1865 On the History of the Last Geological Changes in Scotland.
Quat. Journr. Geol. Soc., 21, pp. 161-203.
- Jamieson, T. F. 1882 On the Causes of the Depression and Re-elevation of the Land During the Glacial Period.
Geol. Mag., 9, pp. 400-407 and 457-466.
- Jardine, W. G. 1963 Pleistocene Sediments at Girvan Ayrshire.
Trans Geol. Soc. Glasgow, 25, pp. 4-16.
- Jardine, W. G. 1964 Post Glacial Sea Levels in South-West Scotland.
Scott. Geogr. Mag., 80, pp. 5-11.
- Johnson, W. A. 1946 Op. cit.
- Jordan, G. F. 1952 Reef Formation in the Gulf of Mexico off Appalachicola.
- Kaye, C. A., and Barghoorn, E. S. 1964 Late Quaternary Sea Level Change and Crustal Rise at Boston, Massachusetts, with Notes on Auto Compaction of Peat.
Bull. Geol. Soc. Am., 75(2), pp. 63-80.
- Keble, R. C. 1947 Notes on Australian Quaternary Climates and Migration.
Mem. Nat. Mus. Victoria (Melbourne), 15, pp. 28-81.
- Keith, M. L., and Degens, E. T. 1959 Geochemical Indicators of Marine and Fresh Water Sediments.
IN Abelson, P. H., editor, Researches in Geochemistry.
N. Y. Wiley, pp. 38-61.
- Kincaid, T. 1957 Local Races and Clines in the Marine Gastropod "Thais Lamellosa" Gonelin, a Population Study.
Calliostoma Co., 75, Seattle.
- King, C. A. M., and Buckley, J. T. 1968 The Analysis of Stone Size and Shape in Arctic Environments.
Journal of Sed. Pet., 38(1), pp. 200-214.
- Kornicker, L. S. Concentration of Ostracods by Alcohol Flotation.

- Kulikov, N., and Martinov, V. 1961 On the Ancient Shorelines of the Bottom of the Kara Sea.
Morskoe Berega Inst. Geol. Acad. Sci. Estonia
S. S. R., Tallin, 8, pp. 147-154.
- Kupfer, D. H. 1964 Five Islands and Mississippi Deltaic Plain Guide Book for Field Strips, South-Eastern Sector.
Geol. Soc. Am., Louisiana, State Univ., pp. 55-64.
- Ladd, H. S., editor, 1957 Treatise on Marine Ecology and Paleocology, V. 2, Paleocology Geol. Soc. Amer. Mem. 67.
- Lakukov, G. I. 1954 The Fluctuation of the Plains of the Polov Basin in the Quaternary Period.
Oceanography 1.
- Laursen, D. 1950 The Stratigraphy of the Marine Quaternary Deposits in West Greenland.
Meddel om Gronland, 151(1).
- Lee, H. A. 1957 Surficial Geology of Frederickton, York, and Sunbury Counties, New Brunswick.
Geol. Survey Canada Paper 56-2.
- Leverett, F., and Taylor, F. B. 1915 The Pleistocene of Indiana and Michigan and the History of the Great Lakes.
U. S. Geol. Survey Monograph 53.
- Lipps, J. H. 1964 Late Pleistocene History of West Anacapa Island, California.
Bull. Geol. Soc. Am., 75(11), pp. 1169-1176.
- Loken, O. 1962 The Late Glacial and Post Glacial Emergence and the Deglaciation of Northernmost Labrador.
Geogr. Bull. 17, pp. 23-56.
- MacGregor, M., and Macgregor, A. G. 1948 The Midland Valley of Scotland.
Geol. Surv. and Mus. Edin. (2nd edit.).
- MacKay, J. R., Matthews, W. H., and MacNeish, R. S. 1961 Geology of the Engigstciak Archaeological Site, Yukon Territory.
Arctic 14(1), pp. 25-52.

- McBurney, C. B. M., and Hey, R. W. 1955 Prehistory and Pleistocene
Geology in Cyrenaican Libya.
Cambridge University Press.
- McCann, S. B. 1961 Some Supposed "Raised Beach" Deposits at
Corran, Loch Linnhe, and Loch Etive.
Geol. Mag. 98, pp. 131-142.
- McCann, S. B. 1963 The Late Glacial Raised Beaches and Readvance
Moraines of the Loch Carran Area, Rossshire.
Scot. Geogr. Mag. 79(3), pp. 164-169.
- McFarlane, P. B. 1955 Survey of Two Drowned River Valleys in Devon.
Geol. Mag. 92(5), pp. 419-429.
- McGill, P. E., and Loranger, D. M. 1961 Micropaleontology.
Foraminifera Zonation.
Geol. of Arctic, Vol. 1, ed. Raasch, pp. 515-531.
- McKee, E. D. 1956 Geology of Kapingamarangi Atoll, Caroline
Islands.
Atoll Research Bulletin 50.
- McKee, E. D. 1958 Geology of Kapinamarangi Atoll, Caroline
Islands.
Bull. Am. Geol. Soc. 69, pp. 241-278.
- McMillan, N. F. 1959 Two Post Glacial Sections at Broighter,
County Londonderry.
Irish Nat. Journ., 13, pp. 16-17.
- McMillan, N. F. 1964 The Mollusca of the Wexford Gravels
(Pleistocene) of South-East Ireland.
Proc. Royal Acad. 63-B(15), pp. 265-289.
- Middleton, G. V., editor 1965 Primary Sedimentary Structures
and Their Hydrodynamic Interpretation.
Soc. Econ. Pal. and Min., Spec. Pub. 12.
- Miller, R. D., and Dobrovolsky, E. 1959 Surficial Geology of
Anchorage and Vicinity, Alaska.
U. S. Geol. Surv. Bull. 1093.
- Mitchell, G. F. 1963 Morainic Ridges on the Floor of the Irish Sea.
Irish Geogr., 4(5), pp. 335-344.
- Murchison 1837

- Nansen, F. 1927 Op. cit.
- Neiheisel, J. 1962 Heavy Mineral Investigation of Recent and Pleistocene Sands of Lower Coastal Plain of Georgia. Bull. Geol. Soc. Am., 73(3), pp. 365-374.
- Newey, W. W. 1966 Pollen Analysis of Sub-Carse Peats of the Forth Valley. Trans. I. B. G., 39, pp. 53-59.
- Nybelin, O. 1946 Nagra Svenska-Subfossilfynd Av Marina Fiskar. Fauna Och Flora, V. 41, pp. 203-211.
- Oaks, R. Q., and Coch, N. K. 1963 Pleistocene Sea Levels, South-East Virginia. Science 140, pp. 979-983.
- Parker, R. H., and Curray, J. R. 1956 Fauna and Bathymetry of Banks on Continental Shelf, North-West Gulf of Mexico. Bull. Am. Ass. Pet. Geol., 40(10), pp. 2428-2439.
- Phleger, F. B. 1951 Ecology of Foraminifera, North-West Gulf of Mexico, Pt. 1, Foraminifera Distribution. Geol. Soc. Amer. Memoir, V. 46, pp. 1-88.
- Phleger, F. B. 1960 Ecology and Distribution of Recent Foraminifera. Johns Hopkins Press, Baltimore.
- Pratt, W. L. 1963 Glauconite from the Sea Floor off Southern California. Unpublished Ph.D., University of California.
- Reid, C. 1913 Op. cit.
- Richards, H. G., and Craig, J. R. 1963 Pleistocene Molluscs from the Continental Shelf off Argentina. Proc. Acad. Nat. Sci., Phila., 115, pp. 127-152.
- Rikhter, V. G., et al 1961 A Contribution to the Study of the Shorelines on the Floor of the Caspian Sea. Doklady Akad. Nauk., U. S. S. R. 135(6), pp. 1476-1479.
- Ritchie, W. 1966 The Post Glacial Rise in Sea Level and Coastal Changes in the Uists. Trans. Inst. Brit. Geog., 39, pp. 79-86.
- Robinson, A. H. W. 1952 The Floor of the British Seas. Scott. Geogr. Mag., 68(2), pp. 64-79.

- Sames, C. W. 1966 Morphometric Dates of Some Recent Pebble Associations and Their Application to Ancient Deposits. Journ. Sed. Pet., V. 36, pp. 126-142.
- Sauramo, M. 1929 The Quaternary Geology of Finland. Comm. Geol. de Finland, Bull. 86, 110 pp.
- Sauramo, M. 1955 Op. cit.
- Schlanger, S. O., and Brookhart, J. W. 1955 Geology and Water Resources of Fanning Island Ulithi Atoll, Western Caroline Islands. Amer. Journ. Sci. 253, pp. 553-573.
- Shepard, F. P. 1956 Late Pleistocene and Recent History of the Central Louisiana Coast. Journ. Geol. 64, pp. 56-69.
- Shepard, F. P. 1960 Rise of Sea Level Along the North-West Gulf of Mexico. IN Recent Sediments North-West Gulf of Mexico. Am. Assoc. Petrol. Geol., pp. 197-220.
- Shepard, F. P. 1964 Sea Level Changes in the Past 6,000 Years. Possible Archaeological Significance. Science 143, pp. 574-576.
- Shepard, F. P., and Moore, D. G. 1965 Central Texas Coast Sedimentation Characteristics of Sedimentary Environment, Recent History and Diagnosis. Bull. Am. Assoc. Pet. Geol. 39(8), pp. 1463-1593.
- Shukri, N. M., and Phillip, G. 1956 The Geology of the Mediterranean Coast Between Rosetta and Bardia, Pt. III, Pleistocene Sediments Mineral Analysis. Bull. Inst. Egypt, 37(2), pp. 445-455.
- Simpson, I. M. 1958 The Pleistocene Succession in the Stockport and South Manchester Area. Quat. Journ. Geol. Soc. Lond., 115(2), pp. 107-121.
- Sissons, J. B. 1962, 1963, 1966 Op. cit.
- Sissons, J. B., and Smith, D. E. 1965 Op. cit.

- Smith, D. E., Sissons, J. B., and Cullingford, R. A. 1969
Isobases for the Main Perth Shoreline in South-East
Scotland as Determined by Trend Surface Analyses.
Trans. Inst. Brit. Geog. 46, pp. 45-55.
- Sjors, H. 1959 Bogs and Fens in the Hudson Bay Lowlands.
Arctic, 14(4), pp. 241-244.
- Sparks, B. W. 1953 Stages in the Evolution of the Weymouth
Lowland.
Trans. Inst. Brit. Geog., 18, pp. 17-29.
- Stainforth, R. M. 1952 Ecology of Arenaceous Foraminifera.
Micropaleontologist, V. 6(1), pp. 42-44.
- Stephens, N. 1958 The Evolution of the Coastline of North-East
Ireland.
Adv. of Sci., V. 16, pp. 389-391.
- Stephens, N. 1963 Op. cit.
- Synge, F. M., and Stephens, N. 1960 The Quaternary Period in
Ireland--An Assessment.
Irish Geog., 4(2), pp. 121-130.
- Stride, A. H. 1959 On the Origin of the Dogger Bank, in the
North Sea.
Geol. Mag., 96(1), pp. 33-44.
- Stride, A. H., and Bowers, R. 1961 Two Unexplored Records of
Quaternary History.
Nature, 191, p. 1001.
- Sugden, J. 1962. The Strand Lines in the Dammen Region of
Alpefjord (Scoresbyland East Greenland).
Geogr. Journ., 128, pp. 43-48.
- Suggate, R. P. 1958 Late Quaternary Deposits of the Christchurch
Metropolitan Area.
N. Z. Jour. Geol. and Geophys. 1, pp. 103-122.
- Te Punga, M. T. 1954 Fossiliferous Late Pleistocene Beds in a
Well at Awchuri, Near Palmerston, North.
N. Z. Jour. Sci. Tech., 36B, pp. 82-92.
- Te Punga, M. T. 1958 Evidence for a Low Sea Level 9,900 Years Ago.
N. Z. Jour. Geol. Geophys. 1(1), pp. 92-94.

- Thomas, R. G. 1963 The Late Pleistocene 150 Foot Fresh Water Beach Line of the Salton Sea Area.
Bull. S. Calif. Acad. Sci., 62(1), pp. 9-17.
- Thorarinsson, S. 1955 Nakudungs Login Vid. Hunafloa--I.
Ljosi Nyrri, Aldursakvardana.
Natturu Fradingurinn, V. 25, pp. 172-186.
- Tinoco I. de M. 1958 Foraminiferos Cuaternarios de Olinda
Estado de Pernambuco.
Civ. de Geol. e Min. Monograph., 14, Rio de Janeiro.
- Upton, J. E. 1954 Terrestrial and Submarine Unconsolidated Deposits in the Vicinity of Eastport, Maine.
Trans. N. Y. Acad. Sci., 16(6), pp. 288-295.
- Valentine, J. W. 1955 Upwelling and Thermally Anomalous Pacific Coast Pleistocene Molluscan Faunas.
Amer. Journ. Sci., V. 253, pp. 463-474.
- Valentine, J. W. 1956 Upper Pleistocene Mollusca from Potrero Canyon, Pacific Palisades, California.
San Diego Soc. Nat. Hist., 12, pp. 181-205.
- Van de Vlerk, I. M. 1950 Correlation Between the Pleistocene Deposits in East Anglia and in the Netherlands.
18th Inter Geol. Cong., London, Pt. 9, pp. 101-106.
- Vedder, J. G., and Norris, R. M. 1963 Geology of San Nicolas Island, California.
U. S. Geol. Survey, Prof. Paper 369.
- Weaver, J. D. 1961 Erosion Surfaces in the Caribbean and Their Significance.
Nature 190(4782), pp. 1186-1187.
- Wenger, S. A. 1951 Elevated Strand Lines of Frobisher Bay, Baffin Bay Island.
Geog. Review, 41, pp. 622-638.
- Wimberley, C. S. 1955 Marine Sediments of Scripps Submarine Canyon, La Jolla, California.
Journ. Sed. Pet., 25(1), pp. 24-37.
- Witting, R. 1918 Hafsyttan Geoidytan och Landhojningen.
Fennia, 39, No. 5, Helsingfors, 346 pp.

Wright, W. B. 1914 The Quaternary Ice Age.

Wyllie, P. J. 1957 A Geological Reconnaissance Through South
Germania Land, North-East Greenland.
Medd om Grønland, 157(1).

CHAPTER 3

BIBLIOGRAPHY

- Agar, R. 1954 Op. cit.
- Agar, R. 1960 Post glacial Erosion of the North Yorkshire Coast
from the Tees Estuary to Ravenscar.
Proc. Yorks. Geol. Soc., V. 32, pp. 409-427.
- Anderson, W. 1939 Possible Late Glacial Sea Levels at 190' and
140' O.D. in the British Isles.
Geol. Mag., 76, pp. 317-321.
- Anderson, W. 1940 Buried Valleys and Late Glacial Drainage
Systems in North-west Durham.
Proc. Geol. Assoc., 51, pp. 274-281.
- Barrow, G. 1888 The Geology of North Cleveland.
Mem. Geol. Survey, England and Wales.
- Beaumont, P. 1967 The Glacial Deposits of Eastern Durham.
Unpublished Ph.D., University of Durham.
- Beaumont, P. 1968 A History of Glacial Research in Northern
England from 1860 to the Present Day.
Dept. Geog., University of Durham, Occas. Paper No. 9,
21 pp.
- Best, R. H. 1954 Westward Proglacial Drainage in Cleveland.
Proc. Yorks. Geol. Soc., V. 30, pp. 301-319.
- Carruthers, R. G. 1939 On Northern Glacial Drifts.
Quat. Journ. Geol. Soc., 95, pp. 299-333.
- Carruthers, R. G. 1946 The Secret of the Glacial Drifts.
Proc. Yorks. Geol. Soc., 27, pp. 43-57.
- Carruthers, R. G. 1953 Glacial Drifts and the Undermelt Theory.
S. Hill, Newcastle.
- Elgee, F. 1908 The Glaciation of North Cleveland.
Proc. Yorks. Geol. Soc., 16, pp. 372-382.
- Elson, J. A. 1956 The Glaciation of the Tiger Hills Region,
Manitoba, Canada.
Unpublished Ph.D., University of Calgary, Alberta,
Canada.

- Fawcett, C. B. 1916 The Middle Tees and Its Tributaries.
Geog. Journ., XLVIII, p. 310.
- Fox, F. 1886 Viaduct Over the River Esk at Whitby.
Min. Proc. instn. Civ. Eng., V. 86, pp. 303-311.
- Gaunt, G. D. 1967 In "Smith and Francis," Geology of the Country
Between Durham and West Hartlepool.
Mem. Geol. Surv.
- Gayner, J. S., and Melmore, S. 1936 Late Glacial Lacustrine
Conditions in the Vale of York and the Tees Basin.
North-West Naturalist, 11, p. 228.
- Goodchild, J. G. 1875 The Glacial Phenomena of the Eden Valley.
Quat. Journ. Geol. Soc., 31, pp. 55-99.
- Harrison, J. W. H. 1921 A Survey of the Lower Tees Marshes and
of the Reclaimed Areas Adjoining Them.
Trans. Nat. Hist. Soc., Northumb., Durham, V. 5, p. 89.
- Kendall, P. F. 1902 A System of Glacier Lakes in the Cleveland
Hills.
Quat. Journ. Geol. Soc., 58, pp. 471-571.
- Lamplugh 1907 (in discussion of Trechmann 1915.)
- Radge, G. W. 1939 The Glaciation of North Cleveland.
Proc. Yorks. Geol. Soc., 24, pp. 180-205.
- Raistrick, A. 1931 The Glaciation of Northumberland and Durham.
Proc. Geol. Assoc., 42, pp. 281-291.
- Raistrick, A. 1934 The Correlation of Glacial Retreat Stages
Across the Pennines.
Proc. Yorks. Geol. Soc., 22, pp. 199-214.
- Smith, D. B., and Francis, E. A. 1967 The Geology of the Country
Between Durham and West Hartlepool.
Mem. Geol. Survey.
- Trechmann, C. T. 1915 Scandinavian Drift on the Durham Coast.
Quat. Journ. Geol. Soc., 71, pp. 53-82.
- Trechmann, C. T. 1919 On a Deposit of Inter-glacial Loess and
Some Transported Pre-glacial Fresh Water Clays on the
Durham Coast.
Quat. Journ. Geol. Soc., 75, pp. 173-203.

CHAPTER 7

BIBLIOGRAPHY

- Aleva, G. J. J. 1956 The Grain Size Distribution of Quartz in
Granitic Rocks of Billiton, Indonesia.
Geologie en Mijnbouw, 18, pp. 177-187.
- Allen, V. T. 1948 Weathering and Heavy Minerals.
Journal of Sed. Pet., 18, pp. 38-42.
- Andel, T. H. van 1959 Reflections on the Interpretation of Heavy
Mineral Analyses.
Journal Sed. Pet., 29, pp. 153-163.
- Andrews, J. T. 1967 Problems in the Analysis of the Vertical
Displacement of Shorelines.
Geog. Bulletin, 9, pp. 71-74.
- Bagnold, R. A. 1954 The Physics of Blown Sand and Desert Dunes.
Methuen, London, 2nd edit., 265 pp.
- Bagnold, R. A. 1966 An Approach to the Sediment Transport
Problem from General Physics.
U. S. Geol. Surv., Prof. paper, 422-I, pp. 11-137.
- Beaumont, P. 1967 The Glacial Deposits of Eastern Durham.
Unpublished Ph.D., University of Durham.
- Beal, M. A., Shepard, F. P. 1956 The Use of Roundness to Determine
Depositional Environments.
Journ. Sed. Pet., V. 26, pp. 49-60.
- Biederman, E. W. 1962 Distinction of Shoreline Environments in
New Jersey.
Journ. Sed. Pet. 32, pp. 181-200.
- Bird, R. J., Freeman, L. O. 1962 Electron Microscopy of Mineral
Particles.
In Milner, H. R., Sedimentary Petrography, V. 1,
pp. 498-521.
- Blatt, H. 1959 Effect of Size and Genetic Quartz Type on Sphericity
and Form of Beach Sediments, Northern New Jersey.
Journ. Sed. Pet., V. 29, pp. 197-206.

- Bradley, J. S. 1957 Differences of Marine and Sub-aerial Sedimentary Environments by Volume Percentage of Heavy Minerals, Mustang Island, Texas.
Journ. Sed. Pet. 27, pp. 116-125.
- Bradshaw, J. S. 1961 Laboratory Experiments on the Ecology of Foraminifera.
Contrib. Cushman Foundation Foram. Res. 12, pp. 87-106.
- Brezina, J. 1963 Kapteyn's Transformation of Grain Size Distribution.
Journ. Sed. Pet. 33, pp. 931-937.
- Briggs, L. I., Middleton, G. V., 1965 Hydromechanical Principles of Sediment Structure Formation.
In Symposium Soc. Econ. Pal. and Min. Spec. Pub. 12, pp. 5-16.
- British Standard 1377 1961 Methods of Testing Soils for Civil Engineering Purposes.
Brit. Standards Institution.
- Bruckner, W. D., Morgan, H. J. 1964 Heavy Minerals on the Continental Shelf of Accra.
Dev. in Sedimentology, Vol. 1, Elsevier, pp. 54-61.
- Bull, W. B. 1962 Relation of Textural (CM) Patterns to the Depositional Environment of Alluvial Fan Deposits.
Journ. Sed. Pet., 32, pp. 211-216.
- Cailleux, A. 1945 Distinction des Galets Marines et Fluviatiles.
Soc. Geol. Bull. France, 5^e Serie, 15, pp. 374-404.
- Carroll, D. 1957 A Statistical Study of Heavy Minerals in Sands of the South River Augusta County, Virginia.
Journ. Sed. Pet. 27, pp. 387-404.
- Chappell, J. 1967 Recognising Fossil Strand lines from Grain Size Analysis.
Journ. Sed. Pet., 37, pp. 157-165.
- Chorley, R. J., Haggett, P. 1965 Trend Surface Mapping in Geographical Research. Trans. Inst. Briti. Geog. 37.
- Currey, Griffiths, 1955 Sphericity and Roundness of Quartz Grains in Sediments.
Bull. Am. Geol. Soc.

- Davis, S. N. 1958 Size Distribution of Rock Types in Stream Gravel and Glacial Till.
Journ. Sed. Pet. 28, pp. 87-94.
- De Graafe, W. P. F. H., Woensdregt, C. F. 1963 Heavy Mineral Analysis of the Beaches Between Ria de Lires and Ria de Murosy Noya (Province La Coruna, N. W. Spain).
Estudios Geologicos, XIX, pp. 9-14.
- Doeglas, D. J. 1946 The Interpretation of the Results of Mechanical Analysis.
Journ. Sed. Pet., 16, pp. 19-40.
- Doeglas, D. J. 1955 An Exponential Function of Size Frequency Distributions of Sediments.
Geol. en Mijnbouw, 18, pp. 1-29.
- Doeglas, D. J. 1962 A Graphic Approach to the Interpretation of Variations in the Size Frequency Distribution of Silty Clays.
Bull. Groupe Franc Argile, 13, pp. 3-14.
- Folk, R. L., Ward, W. C. 1957 Brazos River Bar: A Study in the Significance of Grain Size Parameters.
Journ. Sed. Pet., 27, pp. 3-26.
- Folk, R. L. 1966 A Review of Grain Size Parameters.
Sedimentology, V. 6, pp. 73-94.
- Friedman, G. M. 1961 Distinction Between Dune, Beach, and River Sand from Textural Characteristics.
Journ. Sed. Pet., 31, pp. 514-529.
- Friedman, G. M. 1962 Sorting, Sorting Co-efficients and Log Normality of Grain Size Distribution of Sandstones.
Journ. Geol.
- Friedman, G. M. 1967 Dynamic Processes and Statistical Parameters Compared for Size Frequency Distribution of Beach and River Sands.
Journ. Sed. Pet., 37, pp. 327-354.
- Geer, M. R., Yancey, H. F. 1938 Expression and Interpretation of the Size Composition of Coal.
Trans. A.I.M.E. Coal Div., 130, pp. 250-269.
- Hails, J. R. 1967 Significance of Statistical Parameters Compared for Size Frequency Distribution of Beach and River Sands.
Journ. Sed. Pet., 37, pp. 327-354.

- Hails, J. R., Hoyt, J. H. 1969 The Significance and Limitations of Statistical Parameters for Distinguishing Ancient and Modern Sedimentary Environments of the Lower Georgia Coastal Plain.
Journ. Sed. Pet., 39, pp. 559-580.
- Hand, B. M. 1967 Differentiation of Beach and Dune Sands Using Settling Velocities of Light and Heavy Minerals.
Journ. Sed. Pet., 37, pp. 514-520.
- Harris, S. A. 1957 Mechanical Constitution of Certain Present Day Egyptian Dune Sands.
Journ. Sed. Pet., 27, pp. 421-434.
- Harris, S. A. 1958a Differentiation of Various Egyptian Aeolian Micro Environments by Mechanical Composition.
Journ. Sed. Pet., V. 28, pp. 164-174.
- Harris, S. A. 1958b Probability Curves and the Recognition of Adjustment to Depositional Environment.
Journ. Sed. Pet., 28, pp. 151-163.
- Harris, S. A. 1959 The Mechanical Composition of Some Intertidal Sands.
Journ. Sed. Pet., 29, pp. 414-424.
- Hough, J. L. 1935 The Bottom Sediments of Lake Michigan.
Journ. Sed. Pet., 5, pp. 57-80.
- Imbrie, J. 1963 Factor and Vector Analysis Programs for Analysing Geologic Data.
Office Naval Research, Tech. Rept. 6, Geography Branch, 83 pp.
- Imbrie, J., Van Andel, T. H. 1964 Vector Analysis of Heavy Mineral Data.
Bull. Am. Geol. Soc., 75, pp. 1131-1156.
- Inman, D. L. 1952 Measures for Describing the Size Distribution of Sediments.
Journ. Sed. Pet., 22, pp. 125-145.
- Inman, D. L., Rusnak, G. H. 1956 Changes in Sand Level on the Beach and Shelf at La Jolla, California.
Tech. Memo, B. E. B., 82, 30 pp.

- Kane, W. T., Hubert, J. F. 1963 Fortran Program for Calculation of Grain Size Textural Parameters on the IBM 1620 Computer.
Sedimentology 2, pp. 87-90.
- Kapteyn, J. C. 1903 Skew Frequency Curves.
In P. N. Noordhoff, Groningen, E. Steiger Co., N. Y.
- Kapteyn, J. C. 1916 Skew Frequency Curves in Biology and Statistics.
Recueil des Trav. Bot. Neerl, 13, pp. 105-157.
- King, H. W. 1954 Handbook of Hydraulics.
McGraw-Hill, N. Y., 563 pp.
- King, C. A. M. 1965 Some Observations on the Beaches of the West Coast of County Donegal.
Irish Geography V.
- King, C. A. M., Buckley, J. T. 1968 The Analysis of Stone Size and Shape in Arctic Environments.
Journ. Sed. Pet., 38, pp. 200-214.
- Kirby, R. P. 1968 The Glacial Geomorphology of the Esk Basin, Midlothians.
Unpublished Ph.D. Thesis, University of Edinburgh.
- Klovan, J. E. 1966 The Use of Factor Analysis in Determining Depositional Environments from Grain Size Distributions.
Journ. Sed. Pet., 36, pp. 115-125.
- Klovan, J. E., Billings, G. K. 1967 Classification of Geological Samples by Discriminant Function Analysis.
Bull. Canadian Petroleum Geol., 15, pp. 313-330.
- Krumbein, W. C., Aberdeen, E. J. 1937 The Sediments of Barataria Bay.
Journ. Sed. Pet., 7, pp. 3-17.
- Krumbein, W. C., Pettijohn, F. J. 1938 Manual of Sedimentary Petrography.
D. Appleton-Century Co.
- Krumbein, W. C. 1941 Measurement and Geological Significance of Shape and Roundness of Sedimentary Particles.
Journ. Sed. Pet., 11, pp. 64-72.

- Krumbein, W. C. 1953 Statistical Designs for Sampling Beach Sands.
Trans. Amer. Geophys. Union, 34, pp. 857-868.
- Krumbein, W. C., Slack, H. A. 1956 Relative Efficiency of Beach Sampling Methods.
Beach Erosion Board, Tech. Memo., 90, 43 pp.
- Kuenen, P. H. 1963 Pivotability Studies of Sand in a Shape Sorter.
In Dev. in Sedimentology, I, ed. Van Straatten, pp. 207-215.
- Kuenen, P. H., Humbert, F. L. 1964 Bibliography of Turbidity Currents and Turbidites.
In A. H. Bouma, and A. Brouwer, Turbidites, Elsevier-Amsterdam.
- Leliavsky, S. 1955 An Introduction to Fluvial Hydraulics.
Constable, England, 257 pp.
- Link, A. G. 1967 Delineating the Major Depositional Environments in Northern Port Phillip Bay, Victoria.
Journ. Sed. Pet., 37, pp. 931-951.
- Lockery, A. R. 1967 Report on the Nature and Content of Northumbrian Boulder Clays.
National Coal Board Open Cast Executive, Newcastle, 23 pp.
- MacCarthy, G. R. 1935 Aeolian Sand: A Comparison.
Am. Journ. Sc., 230, pp. 81-95.
- Marshall, P. 1929 Beach Gravels and Sands.
New Zealand Inst. Trans. and Proc., 60, pp. 324-365.
- Martin, W. R. B., Long, A. M. 1960 Heavy Mineral Content and Radio-activity Counts of Beach Sands West of Oreti River Mouth to Blue Cliffs, Southland, New Zealand.
- Martins, L. R. 1965 Significance of Skewness and Kurtosis in Environmental Interpretation.
Journ. Sed. Pet., 35, pp. 768-770.
- Mason, C. C., Folk, R. L. 1958 Differences of Beach, Dune and Aeolian Flat Environments by Size Analysis, Mustang Island, Texas.

- McCammon, R. B. 1962a Moment Measures and the Shape of Size Frequency Distributions.
Journ. of Geol., V. 70, pp. 89-91.
- McCammon, R. B. 1962b Efficiencies of Percentile Measures for Describing the Mean Size and Sorting of Sedimentary Particles.
Journ. of Geol., V. 70, pp. 453-465.
- McKee, E. D. 1966 Structures of Dunes at White Sands National Monument, New Mexico (and a Comparison with Structures of Dunes from other Selected Areas).
Sedimentology, 7, pp. 1-69.
- McKelvey, V. E. 1940 Beach Sediments of Trout Lake, Wisconsin.
Journ. Sed. Pet., 10, pp. 58-64.
- Middleton, G. V. 1965 The Development of Research on Sedimentary Structures.
In Soc. Econ. Pal. and Min. Spec. Pub. 12, pp. 1-4.
- Middleton, G. V. 1965 editor Primary Sedimentary Structures and Their Hydrodynamic Interpretation.
Soc. Econ. Pal. and Min. Spec. Pub. 12.
- Miller, R. L. 1954 A Model for the Analysis of Environments of Sedimentation.
Journ. of Geol., 62, pp. 108-113.
- Miller, R. L. 1956 Trend Surfaces: Their Application to Analysis and Description of Environments of Sedimentation.
Journ. of Geology, V. 64, pp. 425-446.
- Miller, R. L., Zeigler, J. M. 1958 A Study of the Relation Between Dynamics and Sediment Pattern in the Region of Shoaling Waves, Breaker Zone, and Foreshore.
Journ. of Geol., V. 66, pp. 417-441.
- Moiola, R. J., Weiser, D. 1968 Textural Parameters--An Evaluation.
Journ. Sed. Pet., 38, pp. 45-53.
- Moss, A. J. 1962 The Physical Nature of Common Sandy and Pebbly Deposits. Part I.
Am. Journ. Sci., 260, pp. 337-373.
- Moss, A. J. 1963 The Physical Nature of Common Sandy and Pebbly Deposits. Part II.
Am. Journ. Sc., 261, pp. 297-343.

- Mueller, G. 1967 Methods in Sedimentary Petrology.
Hafner Pub. Co., N. Y., 283 pp.
- Passega, R. 1957 Texture as a Characteristic of Elastic Deposition.
Am. Assoc. Petrol. Geol. Bull. 41, pp. 1952-1984.
- Peikert, E. W. 1963 IBM 709. Program for Least Squares Analysis
of Three-Dimensional Geological and Geophysical
Observations.
Tech. Rep. 4, O. N. R., No. 389-135.
- Plumley, W. J., Davis, D. H. 1956 Estimation of Recent Sediment
Size Parameters from a Triangular Diagram.
Journ. Sed. Pet., V. 26, pp. 140-155.
- Pollack, J. M. 1961 Significance of Compositional and Textural
Properties of South Canadian River Channel Sands,
New Mexico, Texas, and Oklahoma.
Journ. Sed. Pet., 31, pp. 15-37.
- Poole, D. M. 1958 Heavy Mineral Variation in San Antonio and
Mesquite Bays of the Central Texas Coast.
Journ. Sed. Pet., 28, pp. 65-74.
- Porter, J. J. 1962 Electron Microscopy of Sand Surface Texture.
Journ. Sed. Pet., 32, pp. 124-135.
- Potter, P. E., Pettijohn, F. J. 1963 Paleocurrents and Basin
Analysis.
New York. Academic Press Inc., 296 pp.
- Pratt, W. L. 1962 The Origin and Distribution of Glauconite from
the Sea Floor off California and Baja, California.
Unpublished Ph.D., University of Southern California.
- Reineck, H. E. 1967 Vergleich von Schlamm-Analysen mit der
Pipette Nach Kohn und dem Araometernach Casagraade.
Senckenbergia Leth., 48, pp. 351-357.
- Rogers, J. J. W., Powell, W. F. 1958 Size Distributions of
Zircon Grains.
Journ. Sed. Pet., 28, pp. 36-39.
- Rogers, J. J. W., Adams, H. C. jnr. The Mineralogy and Texture of
Beach Sands of Galveston Island, Texas.
Journ. Sed. Pet., 29, pp. 207-211.

- Rosin, P., Remmler, E. 1933 The Laws Governing the Fineness of Powdered Coal.
Journ. Inst. Fuel, 7, pp. 29-36.
- Ruhkin, L. B. 1937 A New Method for the Determination of the Conditions of Deposition of Ancient Sands.
Pro. Sov. Geol., 7, pp. 953-959.
- Sahu, B. K. 1964 Depositional Mechanisms from the Size Analysis of Elastic Sediments.
Journ. Sed. Pet., 34, pp. 73-83.
- Sames, C. W. 1966 Morphometric Data of Some Recent Pebble Associations and Their Application to Ancient Deposits.
Journ. Sed. Pet., 36, pp. 126-142.
- Sevon, W. D. 1966 Distinction of New Zealand Beach, Dune and River Sands by their Grain Size Distribution Characteristics.
N. Z. Journ. Geol. and Geophys., 9, pp. 212-223.
- Sevon, W. D. 1967 First and Second Degree Regression Correlation of the Size Analysis Statistical Parameters of Trask, Inman, Folk and Ward, and Friedman.
Journ. Sed. Pet., 37, pp. 238-240.
- Shepard, F. P., Moore, D. G. 1955 Central Texas Coast Sedimentation: Characteristics of Sedimentary Environment, Recent History and Diagenesis.
Bull. Am. Assoc. Petrol. Geol., 39, pp. 1463-1593.
- Shepard, F. P., Young, R. 1961 Distinguishing Between Beach and Dune Sands.
Journ. Sed. Pet., 31, pp. 196-214.
- Shepard, F. P. 1964 Criteria in Modern Sediments Useful in Recognising Ancient Sedimentary Environments.
Devs. in Sedimentology, Vol. 1, Elsevier, pp. 1-25.
- Smalley, I. J. 1967 The Presentation of Subjective Shape and Roundness Data.
Sedimentology, Vol. 8, pp. 35-38.
- Sneed, E. D., Folk, R. L. 1958 Pebbles in the Lower Colorado River, Texas. A Study in Particle Morphogenesis.
Journ. Geol., V. 66, pp. 114-150.

- Solohub, J. T. 1967 Grand Beach. A Test of Grain Size Distribution Statistics as Indicators of Depositional Environments. Unpub. M.Sc. thesis, University of Manitoba, Canada.
- Spencer, D. W. 1963 The Interpretation of Grain Size Distribution Curves of Elastic Sediments. Journ. Sed. Pet., 33, pp. 180-190.
- Stairmond, C. J. 1947 Symposium on Particle Size Analysis. Supplement to Trans. Inst. Chem. Engineers, V. 25, p. 128.
- Tanner, W. F. 1958 The Zig-Zag Nature of Type I and Type IV Curves. Journ. Sed. Pet., 28, pp. 372-375.
- Tiechert, C. 1958 Cold and Deep Water Coral Banks. Bull. Am. Assoc. Petrol. Geol., 42, pp. 1064-1082.
- Thurstone, L. L. 1947 Multiple Factor Analysis. University of Chicago Press, 535 pp.
- Udden, J. 1898 Mechanical Composition of Wind Deposits. Augstan Library Publication, 1, 69 pp.
- Udden, J. 1914 Mechanical Composition of Elastic Sediments. Geol. Soc. Am. Bull., 25, pp. 655-744.
- Van Straaten, L. M. J. U. 1954 Sedimentology of Recent Tidal Flat Deposits and the Psammites du Condroz. Geol. en Mijn., N. S. 16, pp. 25-47.
- Van Straaten, L. M. J. U., editor 1964 Deltaic and Shallow Marine Deposits. Elsevier. Development in Sedimentology, Vol. 1.
- Visher, G. S. 1965 Fluvial Processes as Interpreted from Ancient and Recent Fluvial Deposits. In G. V. Middleton, editor, Primary Sedimentary Structures and Their Hydrodynamic Interpretation. Soc. Econ. Pal. and Min., Spec. Publ. 12, pp. 116-132.
- Wadell, H. 1932 Volume, Shape, and Roundness of Rock Particles. Journ. Geol., Vol. XL, pp. 443-451.

- Waskom, J. D. 1958 Roundness as an Indicator of Environment
Along the Coast of Panhandle, Florida.
Journ. Sed. Pet., V. 28, pp. 351-360.
- White, J. R., Williams, E. G. 1967 The Nature of a Fluvial
Process as Defined by Settling Velocities of Heavy
and Light Minerals.
Journ. Sed. Pet., 37, pp. 530-539.
- Whitten, E. H. T. 1959 Composition Trends in Granite, Modal
Variation, and Ghost-Stratigraphy in Part of the
Donegal Granite, Eire.
Journ. Geophys. Research, V. 64, pp. 835-848.
- Whitten, E. H. T. 1961 Quantitative Areal Modal Analysis of
Granitic Complexes.
Geol. Soc. Amer. Bull., V. 72, pp. 1331-1360.
- Williams, E. M. 1965 A Method of Indicating Pebble Shape with
One Parameter.
Journ. Sed. Pet., 35, pp. 993-996.
- Zingg, T. 1935 Beitrag Zur Schotteranalyse.
Schweizer Mines. Petrog Mitt., 15, pp. 39-140.

CHAPTER 8BIBLIOGRAPHY

- Apfel, E. T. 1938 Phase Sampling of Sediments.
Journal of Sedimentary Petrology, V. 8, pp. 67-68.
- Beaumont, P. 1967 Op. cit. (Chapter 3)
- Chappell, J. 1967 Op. cit. (Chapter 7)
- Cochran, W. G., Mosteller, F., Tukey, J. W. 1954 Principles of Sampling.
Journ. Am. Stat. Soc., p. 13.
- Emery, K. O., and Stevenson, R. E. 1950 Laminated Beach Sand.
Journ. Sed. Pet., V. 20, pp. 220-223.
- Gees, R. A. 1969 Sampling of Laminated Beach Sands.
Maritime Sediments, V. 5, No. 2, pp. 40-43.
- Griffiths, J. C. 1967 Scientific Methods in Analysis of Sediments.
McGraw-Hill, New York.
- Hails, J. 1967 Op. cit. (Chapter 7)
- Harris, S. A. 1959 Op. cit. (Chapter 7)
- Klovan, J. E., Solohub, J. 1970 Op. cit. (Chapter 7)
- Krumbein, W. C. 1953 Statistical Designs for Sampling Beach Sand.
Am. Geophys. Union Trans., V. 34, pp. 857-868.
- Krumbein, W. C., Slack, H. A. 1956 Relative Efficiency of Beach Sampling Methods.
Tech. Memo 90, B. E. B. Corps Engineers, 52 pp.
- McIntyre, D. D. 1959 The Hydraulic Equivalence and Size Distributions of Some Mineral Grains from a Beach.
Journ. Geol., V. 67, pp. 278-301.
- Middleton, G. V. 1965 Op. cit. (Chapter 7)
- Moiola, R. J., Weiser, D. 1968 Op. cit. (Chapter 7)

- Otto, G. H. 1938 The Sedimentation Unit and its Use in Field Sampling.
Journ. Geol., V. 46, pp. 569-582.
- Ruhkin, L. B. 1937 A New Method for the Determination of the Conditions of Deposition of Ancient Sands.
Prob. Sov. Geol., V. 7, pp. 953-959.
- Sevon, W. 1965 Op. cit. (Chapter 7)
- Robinson, A. H. W. 1968 The Submerged Glacial Landscape off the Lincolnshire Coast.
Inst. Brit. Geog. Trans., V. 44, pp. 119-132.
- Solohub, J. 1967 Op. cit. (Chapter 7)
- Thompson, W. O. 1937 Original Structures of Beaches, Bars, and Dunes.
Geol. Soc. Am. Bull., V. 48, pp. 723-752.
- Udden, J. 1914 Op. cit. (Chapter 7)

Chapter Ten

Bibliography

- Agar R. 1954 op. cit. Chapter 3
- Anderson W. 1939 op. cit. Chapter 3
- Anderson, W. 1947 Discussion of Trechmann 1947
op. cit.
- Barrow 1888 op. cit. Chapter 3
- Beaumont P. 1967 op. cit.
- Beaumont P. 1970 in press British Association for the
Advancement of Science Pamphlet.
Durham meeting Sept. 1970.
- Carruthers R. G. 1953 op. cit. Chapter 3
- Dietz R. S. 1942 Clay minerals in recent marine
sediments
Am. Min. V 27 p. 219-220.
- Gaunt G. D. 1967 (In Smith and Frances 1967) op. cit. Chapter 3
- Jorgensen P. 1965 Mineralogical composition and
weathering of some late
pleistocene marine clays from
the Kongsvinger area. S. Norway.
Geologiska. Foreningens.
Stockholm V 87 pt. 1 pp.62-83
- Powers M.C. 1957 Adjustment of land derived clays
to the marine environment
Journ. Sed. Pet. V 27 pp. 355
- 372.
- Pratt W. L. 1962 op. cit. Chapter 7
- Smith D. B. Francis E. A. 1967 op. cit. Chapter 3
- Tute J. S. 1883 op. cit. Chapter 3
- Veitch W. Y. 1883 op. cit. Chapter 3

Chapter Eleven


Bibliography

- Agar R. 1954 op. cit. Chapter 3.
- Barrow G. 1888 op. cit. Chapter 3.
- Blackburn K. B. 1952. The dating of a deposit contain-
ing an Elk skeleton found at
Neasham near Darlington, Co.
Durham.
New Phytologist V51 P 364-377
- Cameron A. G. 1878 Notes on some peat at Kildale
and West Hartlepool
Geol Mag. V5 P 351
- Gadd N. R. 1964 Moraines in the Appalachian
region of Quebec. Bull Geol
Soc. Amer. V75 (2) PP 1249-1254.
- Ingram J. 1909 A note on the Peat deposit at
Hartlepool.
Proc. Cleveland Nat. Field Club
VIII P 217
- Jelgersma S. 1961 op. cit. Chapter one.
- Radge G. W. 1939 op. cit. Chapter 3.
- Simpson H. 1904 Note on the submerged forest
peat beds at Redcar.
Proc. Cleveland Nat. Field Club
Vol. I P 274
- Smith and Gaunt 1967 (in Smith and Francis 1967 op.
cit.) Chapter 3.
- Trechmann C. T. 1947 op. cit. Chapter 3.



PHOTOGRAPHS.

PLATE I.



Supposed raised marine sand. Cat Nab, Saltburn.

PLATE II.




BREAK OF SLOPE NEAR HOWDEN HALL FARM. LOWER TEES.

PLATE III.




SUPPOSED RAISED STORM BEACH DEPOSITS NEAR HARTLEPOOL.

PLATE IV.



SEA BED GRAB SAMPLER AS USED IN THE TEES BAY.

PLATE V.



SEA BED BUCKET DREDGE AS USED IN THE TEES BAY.


MARINE SURVEY.

PLATE VI. VESSEL, EQUIPMENT, AND PERSONNEL USED DURING
SAMPLING AND SOUNDING CRUISES.

PLATE VII. ECHO SOUNDER TRANSDUCER.

PLATE VIII. ECHO SOUNDING CHART RECORDER.







APPENDICES.

APPENDICES.

APPENDIX INATIONAL GRID REFERENCES OF PLACE NAMES REFERREDTO IN THE TEXT

Names are listed in the order they occur in the text.

O.S., Sheets 84, 85, 86.

THE LOWER TEES

<u>Place Name</u>	<u>Location</u>
Cleveland Hills	585017
Egglescliffe	421135
Billingham Beck	436233
River Leven	449114
Ormesby Beck	511181
Middle Beck	495175
Stainmore Pass	914109
Hutton Henry	425361
Stokesley	520084
Warrenby	581250
Coatham	591252
Redcar	615249
Saltburn	667215
Whitby	905109
Easington	417435
Hartlepool	522333
Throston	491338
Greatham	492277
Thorpe Thewles	400233

The Lower Tees - continued

<u>Place Name</u>	<u>Location</u>
Viewly Hill Farm	435237
Stockton	445225
Skelton Beck	667215
Windy Hill Farm	652218
Marske	645230
Ox Close Farm	651211
Pontac Farm	626211
Fell Briggs	613210
Sparrow Park Farm	621211
Turners House	601210
Long Beck House	625217
Cat Flats	623217
Black's Bridge	621215
Wheat Acres Farm	599218
Yearby	600210
Middlesbrough	495186
Kirkleatham Hall	595209
Fox Rush Farm	587221
Broadway Farm	560205
Town Farm	582207
Lackenby Hall	565195
Normanby High Farm	535195
Spencer Beck Farm	540180
Coulby Manor	496154
Sandy Flat Farm	493158
Stainsby Hall Farm	471152
High Leven	450123
Maltby	466137

The Lower Tees - continued

<u>Place Name</u>	<u>Location</u>
Barwick Farm	432146
Thornaby Vale	455145
Leven Month Farm	435122
Morley Carr	412110
Spittal Flat	428117
Howe Hill Cottages	355110
Hill House	354102
Over Dinsdale	362100
Girsby Grange Farm	385086
Old School (Girsby)	358094
Low Entercommon	335060
Eryholme	321088
Low Hail Farm	309097
Breakhouse Bank	332084
Garden House Farm	303106
Hurworth	305104
Hurworth Moor	315121
Middleton St. George	355140
Long Newton	380162
Tees Airport	375130
Elton	401174
Betty's Farm	405190
Coatham Stob	405162
Howden Hall	415225
Warren House Farm	437254
Woodside	436273
Wolviston	454258
Middle Burn Toft	457279

The Lower Tees - continued

<u>Place Name</u>	<u>Location</u>
Marsh House	461256
High Grange	457251
North Burn	462274
Claxton	476287
Cowpen Bewley	483247
Throston Grange	492341
Middle Warren	494346
Springwell House	488355
I.C.I. Wilton	585198
Girsby	355085
Seaham	437493
South Gare	557277
North Gare	542284
Seaton Carew	513301
Wynyard	418250
Stob House Farm	457272
Brierton	476300
Crimdon	487365
Cat Nab	668216
Huntcliff	691219
Newton Bewley	475266
Dalton Piercy	465312
Dinsdale	330115
Neasham	331102
The Slake	515328
Seal Sands	525252

APPENDIX IILEVELLED HEIGHTS OF RAISED TERRACES

<u>Terrace</u>	<u>Height (in feet)</u>	<u>Location</u>
Skelton Beck, Saltburn	13.60	668217
" " "	13.35	
" " "	16.36	
" " "	17.26	
" " "	19.25	667213
Windy Hill	134.42	652218
" "	133.98	
" "	133.67	
" "	133.41	
" "	133.52	
" "	132.78	
" "	132.43	
" "	132.19	
" "	132.18	650219
Ox Close Farm	141.2	651211
" " "	143.16	
" " "	151.69	
" " "	152.11	
" " "	154.23	
" " "	155.07	
" " "	156.43	
" " "	154.12	
" " "	162.81	
" " "	164.45	
" " "	169.72	
" " "	170.08	
" " "	168.48	

<u>Terrace</u>	<u>Height (in feet)</u>	<u>Location</u>
Pontac Farm	168.01	626211
Fell Briggs	151.40	613210
" "	155.82	
" "	157.15	
" "	157.44	
Sparrow Park Farm	157.67	621211
Turners House	84.90	601210
" "	85.50	
" "	84.74	
" "	83.31	
" "	84.38	
" "	84.40	
" "	85.64	
" "	85.47	
" "	83.45	
" "	84.83	
" "	84.70	
" "	84.68	
Long Beck House	84.82	625217
Blacks Bridge	47.43	621215
" "	47.28	
" "	47.14	
" "	46.97	
" "	47.39	
" "	46.98	

<u>Terrace</u>	<u>Height (in feet)</u>	<u>Location</u>
Wheat Acres Farm	46.54	599218
Kirkleatham Hall	55.65	595207
" "	55.23	
" "	54.61	
" "	54.71	
" "	55.14	
" "	55.67	585215
ICI Wilton	52.56	580210
" "	52.91	
" "	51.91	
" "	52.50	
" "	52.47	
" "	52.42	
" "	52.37	575209
Broadway Farm	80.44	560205
" "	81.20	
" "	80.47	
" "	81.13	
" "	80.78	
" "	80.91	
" "	81.37	
" "	81.55	
Broadway Farm	78.92	560206
" "	78.76	
" "	78.48	

<u>Terrace</u>	<u>Height (in feet)</u>	<u>Location</u>
Broadway Farm	93.11	577203
" "	93.12	
" "	93.08	
" "	92.97	
" "	92.68	
" "	92.31	
" "	92.30	
Town Farm	92.17	582207
Lackenby Hall	75.68	565195
" "	75.79	
" "	76.31	
" "	77.09	
" "	77.18	
Spencer Beck Farm	100.07	540180
" " "	100.77	
" " "	98.59	
" " "	100.13	
" " "	100.38	536177
Normanby High Farm	50.69	535195
" " "	50.32	
" " "	49.91	
" " "	49.65	
" " "	49.42	
" " "	49.38	534195

<u>Terrace</u>			<u>Height (in feet)</u>		<u>Location</u>
Coulby Manor			<u>High Terrace</u>	<u>Low Terrace</u>	496154
"	"		108.22	106.96	
"	"		109.95	103.72	
"	"		106.21	105.73	
"	"		108.12	105.51	
"	"		107.89	104.24	
"	"			105.26	494154
Sandy Flats Farm			99.63	Low Terrace	493158
"	"	"	99.37	46.73	
"	"	"	99.14	46.43	
"	"	"	99.01		
"	"	"	98.94		
"	"	"	98.87		
"	"	"	98.93		497158
Stainsby Hall Farm			77.93		471152
"	"	"	77.40		
"	"	"	77.18		
"	"	"	77.29		
"	"	"	77.28		468152
High Leven			122.24	116.75	450123
"	"		122.22	116.41	
"	"		122.13	115.77	
Maltby			122.05	115.33	466137
"				115.30	

<u>Terrace</u>		<u>Height (in feet)</u>	<u>Location</u>
Barwick Farm		76.71	432146
Leven Mouth Farm		118.39	435122
" " "		118.40	
" " "		118.36	
" " "		118.21	
" " "		119.0	
" " "		118.61	
" " "		118.37	
" " "		118.35	
" " "		118.30	
" " "		118.31	
Morley Carr		118.31	412110
Howe Hill Cottages	1)	77.13	355110
" " "		77.77	
" " "		78.67	
" " "		79.04	
" " "		79.20	
" " "		79.50	
" " "		80.37	
" " "		80.45	
Howe Hill Cottages	2)	108.12	355110
" " "		106.78	
Howe Hill Cottages	3)	113.39	
" " "		113.95	
" " "		113.56	

<u>Terrace</u>	<u>Height (in feet)</u>	<u>Location</u>
East of Hill House	121.32	362100
" "	122.49	
" "	122.31	
" "	122.84	
" "	120.34	
" "	118.49	
" "	120.75	
" "	121.97	
" "	122.60	
" "	122.78	
Rose Hill Farm	123.23	358104
Girsby Grange	144.76	385086
" "	144.78	
" "	144.89	
" "	144.92	
" "	144.95	
" "	145.02	
" "	145.02	
Old School Girsby	1) 132.4	358094
" "	135.16	
" "	132.26	
" "	135.22	
" "	135.37	
" "		
" "	2) 128.44	
" "	129.42	
" "	128.83	
" "	129.08	
" "	129.50	
" "	129.59	

<u>Terrace</u>		<u>Height (in feet)</u>	<u>Location</u>
Low Entercommon	1)	174.24	335060
" "		174.55	
" "		174.61	
" "		174.72	
" "	2)	166.13	335060
" "		166.24	
" "		166.41	
Low Hail Farm		134.93	309097
" "		134.40	
" "		134.73	
" "		134.62	
" "		134.68	
" "		134.62	
" "		134.49	
" "		134.85	
Garden House Farm, Hurworth		118.20	303106
" "	"	120.22	
" "	"	119.60	
" "	"	120.22	
" "	"	118.05	
" "	"	118.05	
Hurworth Moor		131.67	315121
" "		130.94	
" "		130.82	
" "		131.30	
" "		131.60	
" "		130.75	

<u>Terrace</u>	<u>Height (in feet)</u>	<u>Location</u>
Middleton St. George	132.82	355140
" "	130.82	
" "	136.69	
" "	130.91	
" "	131.60	
" "	130.73	
" "	131.12	
" "	134.57	
" "	134.78	
" "	135.87	
" "	135.55	
" "	137.55	
" "	137.45	
Long Newton	137.84	380162
Long Newton	113.08	380162
" "	116.17	
" "	115.89	
" "	113.19	
" "	115.45	
" "	117.56	
" "	119.00	
" "	117.40	
" "	116.38	
" "	116.58	
Middleton St. George	119.47	355140

<u>Terrace</u>		<u>Height (in feet.)</u>	<u>Location</u>
Elton		105.73	401174
"		105.77	
"		104.78	
"		104.67	
"		105.39	
"		105.13	
"		105.76	
Betty's Farm		106.42	405190
Coatham Stob	1)	94.81	
" "		94.82	
" "			
" "	2)	79.75	
" "		79.82	
" "		79.94	
" "		80.21	
" "		80.29	
Coatham Stob (sand layer)	1)	94.86	405162
" "		94.86	
" "		94.87	
" "		94.88	
" "		94.89	
" "		94.89	
" "	2)	95.33	
" "		95.50	
" "		95.43	
" "		95.47	
" "		95.54	
" "		95.69	

<u>Terrace</u>	<u>Height (in feet)</u>	<u>Location</u>
North edge of Stockton	59.58	445225
	59.63	
Howden Hall	131.96	415225
" "	132.20	
" "	132.95	
" "	136.24	
Warren House Farm	111.36	437254
" " "	111.40	
" " "	112.06	
" " "	111.98	
" " "	112.01	
" " "	112.23	
" " "	112.49	
" " "	112.59	
Woodside	176.79	436273
"	175.94	
"	176.67	
"	178.80	
"	177.74	
Wolviston	109.91	454258
"	110.36	
"	110.73	
"	110.30	
"	109.63	
"	109.44	
"	107.42	

<u>Terrace</u>	<u>Height (in feet)</u>		<u>Location</u>
Wolvison (continued)	106.93		
"	108.56		
"	106.07		
"	108.51		
"	110.68		
Middle Burn Toft	109.94		457279
Marsh House	1)	59.81	2) 59.75
" "		59.78	59.51
" "		59.78	59.26
" "		59.66	59.25
" "		59.57	
" "		59.51	
High Grange	59.23		457251
Middle Burn Toft	1) 53.55	2) 73.75	3) 68.45 457279
" " "	54.01	73.61	68.45
" " "	53.14	73.54	68.41
" " "	53.02	73.44	67.55
" " "	52.44	73.32	67.45
" " "			67.31
" " "			66.48
Claxton	61.24		476287
"	61.13		
"	61.30		
"	61.61		

<u>Terrace</u>	<u>Height (in feet)</u>		<u>Location</u>
Claxton	1) 67.97	2) 45.19	476287
"	69.17		
"	67.38		
"	66.02		
Cowpen Bewley	19.05		483247
" "	17.51		
" "	15.82		
" "	14.80		
" "	14.64		
Throston Grange	60.19		492341
" "	60.23		
" "	58.33		
" "	59.32		
" "	60.34		
" "	60.41		
" "	60.52		
" "	60.60		
" "	60.22		
" "	60.99		
Middle Warren	60.66		494346

APPENDIX 3

METHODS OF SEDIMENTARY ANALYSES

1) Particle Size Analysis - Dry Sieving

- 1) Encompasses Size Range -2ϕ to $+4.2\phi$ (coarse sand to fine sand).

2) Apparatus

- (a) British Standard Sieves plus shaker.
- (b) Balance accurate to 0.01 grams.
- (c) Sample Riffler.
- (d) Drying Oven (constant temperature 105° to 110°C).
- (e) Mortar and Pestle.
- (f) Evaporating dishes.
- (g) Spatulas.

3) Procedure

- (a) Sample dried.
- (b) Sample riffled to obtain 100 gram sample.
- (c) Sample checked for grain conglomerates which result from inadequate washing before drying. (Only applicable to salt water samples).
- (d) Sample sieved for up to 15 minutes.
- (e) Residue on each sieve weighed and recorded as a percentage of the total sample weight
- (f) Results recorded and presented graphically on arithmetic probability paper.

HEAVY MINERAL ANALYSIS -(See accompanying table)

Procedure

- (a) Sample residues from the British Standard Sieves 72, 100, 150, 200, rewashed to remove dust.
- (b) Sample redried.
- (c) Sample subjected to hand magnet to remove magnetic and pyrrhotite.
- (d) Sample put through S.G. Franz Isodynamic separator (electro magnet.) (At least twice if not more times to obtain best separation.
- (e) Weight of sample recorded before and after processing to determine weight loss.
- (f) Weight loss recorded as a percentage of sample size fraction.

Other Mineralogical Studies

1) Mica Content

- (a) Microscopic grain count per 1000 on each of the above four size fractions. (72, 100, 150, 200.)

Best results were obtained using X10 magnification under reflected light, with the grains on a black matt finished base.

2) Coal Content

(As above but with white background.)

3) Shell Content

(As above with black background.)

Other Mineralogical Studies - continued

4) Rock Fragments

(As above with black background and X20 magnification on smaller size fractions.)

APPENDIX 4*PARTICLE SIZE ANALYSISSamples Used for Factor Analysis 1

Sieve Size	Percentage								
British Standard	Sample Number								
	W1	W2	W3	W4	W5	W6	W7	W8	W9
8	0.04	0.04	-	-	0.08	0.08	2.92	0.08	0.2
14	0.02	0.06	0.14	0.12	0.08	0.48	2.41	0.42	0.82
25	0.36	0.8	0.34	0.38	1.21	1.29	4.08	1.29	3.02
36	0.64	1.04	0.56	0.58	0.4	0.58	1.91	0.78	5.61
52	1.5	2.02	0.66	0.76	0.78	1.01	2.91	2.1	24.16
72	2.4	3.0	1.2	1.2	2.8	4.92	9.31	7.19	46.23
100	5.5	6.26	3.6	3.8	23.42	37.77	31.54	10.75	16.51
150	37.9	37.90	67.2	66.4	53.57	47.8	28.88	33.0	2.2
200	26.8	28.22	20.6	21.06	8.95	3.89	4.31	17.2	0.14
240	12.56	11.06	2.86	3.08	2.26	0.51	3.4	9.2	0.07

Sieve Size	Percentage								
	Sample Number								
	W10	W11	W12	W13	W14	W15	W16	W17	W18
8	0.02	-	0.06	0.2	0.08	0.1	0.05	0.1	0.1
14	0.26	0.2	0.39	0.45	0.3	0.4	0.29	0.2	0.18
25	2.26	1.68	1.95	2.6	1.1	1.2	1.28	0.85	0.92
36	4.89	0.7	1.12	2.89	1.22	1.15	1.59	0.3	0.8
52	10.29	1.48	2.48	13.49	5.35	5.21	5.9	0.53	0.8
72	41.85	4.18	6.2	46.15	17.68	18.1	18.93	0.62	1.08
100	27.23	7.2	8.0	23.91	19.78	19.07	19.2	6.1	6.4
150	4.33	30.7	31.07	4.09	34.28	35.98	34.64	38.42	40.9
200	0.11	21.0	18.5	0.1	12.91	12.1	11.26	33.5	28.1
240	0.01	11.12	10.39	0.01	3.62	3.47	3.08	12.28	12.9

* Appendices 4 and 5 have been restricted primarily to those samples which were used for factor analysis. This was done because many of the results were included in the text in graphical or summary table form and also because it was factor analysis which provided the key to the differentiation of depositional environments.

Sieve Size	Percentage								
British Standard	Sample Number								
	W19	W20	W21	W22	W23	W24	W25	W26	W27
8	1.1	0.27	0.4	0.08	0.51	0.27	0.38	0.09	0.09
14	1.82	0.41	0.71	0.2	0.5	0.52	0.35	0.3	0.21
25	3.16	1.75	2.11	1.08	0.92	1.3	1.01	0.95	0.95
36	1.28	0.99	0.91	0.81	0.32	0.48	0.48	0.42	0.44
52	2.22	1.09	1.02	0.92	0.6	0.7	0.78	0.71	0.69
72	8.0	2.0	2.5	1.7	2.95	2.8	3.59	3.59	4.03
100	29.7	12.48	16.17	10.92	19.58	18.61	23.67	24.03	25.38
150	46.5	36.09	36.8	34.1	54.09	55.79	53.33	56.25	55.37
200	4.8	15.89	14.0	17.1	16.21	15.3	12.98	11.2	10.5
240	0.44	8.09	6.51	8.03	2.76	2.09	1.68	1.3	1.13

Sieve Size	Percentage								
	Sample Number								
	W28	W29	W30	W31	W32	W33	W34	W35	W36
8	0.05	0.93	1.44	0.5	1.14	0.01	0.02	0.89	13.18
14	0.14	0.7	1.71	0.85	0.83	0.03	0.11	1.41	6.11
25	0.62	2.63	2.52	2.45	2.05	0.53	0.8	3.7	13.93
36	0.38	1.44	1.1	1.6	1.32	0.38	0.4	1.0	9.89
52	0.6	3.75	2.98	4.02	3.6	0.61	0.62	1.08	22.63
72	3.98	25.33	21.1	24.92	24.52	1.8	2.25	2.4	26.25
100	27.38	53.92	54.9	52.55	52.6	14.53	16.13	15.3	6.52
150	53.5	9.95	12.58	11.6	12.22	48.63	48.79	42.99	0.74
200	11.02	0.6	0.92	0.7	0.8	19.05	16.31	13.89	0.05
240	1.1	0.1	0.15	0.08	0.12	6.54	5.91	5.75	-

Sieve Size	Percentage								
British Standard	Sample Number								
	W37	W38	W39	W40	W41	W42	W43	W44	W45
8	0.4	0.6	3.6	1.1	2.0	1.81	2.3	60.9	57.6
14	1.12	18.6	3.02	1.1	1.51	1.23	1.3	10.7	9.7
25	2.6	24.42	7.6	1.9	1.89	2.08	1.13	11.92	12.89
36	2.6	10.41	5.9	0.98	0.81	0.98	0.8	3.32	4.29
52	14.41	10.52	15.28	1.7	1.61	2.01	1.86	2.6	3.5
72	48.23	9.0	36.46	13.6	13.9	13.47	15.32	2.0	3.0
100	23.2	6.51	18.5	58.4	59.55	57.89	59.59	5.7	5.78
150	4.18	5.97	5.91	19.82	17.37	19.73	15.9	2.21	2.21
200	0.72	3.21	1.18	0.8	1.1	0.7	0.58	0.14	0.27
240	0.38	1.52	0.42	0.1	0.11	0.1	0.09	0.08	0.11

Sieve Size	Percentage								
	Sample Number								
	W46	W47	W48	W49	W50	W51	W52	W53	W54
8	70.2	0.18	0.09	0.15	0.19	0.13	0.08	0.41	0.21
14	11.47	0.25	0.58	0.6	0.47	0.49	0.1	0.8	0.64
25	11.16	0.8	3.03	3.48	1.95	1.59	1.38	2.21	1.31
36	2.8	0.62	4.42	3.56	2.25	1.5	0.68	1.2	0.61
52	1.71	1.95	12.0	9.23	7.45	4.82	1.47	1.81	1.17
72	0.75	17.37	23.5	22.78	23.9	19.0	6.85	12.1	6.42
100	1.13	69.1	32.03	33.1	37.13	40.5	65.47	60.02	62.87
150	0.47	9.38	20.85	23.81	23.61	28.9	22.6	19.64	25.03
200	0.09	0.12	1.69	1.75	1.58	1.7	0.7	0.76	0.97
240	0.03	0.05	0.2	0.18	0.11	0.2	0.11	0.11	0.19

Sieve Size	Percentage								
British Standard	Sample Number								
	W55	W56	W57	W58	W59	W60	W61	W62	W63
8	0.03	3.0	3.3	0.15	0.1	0.1	0.1	0.1	0.15
14	0.39	1.88	2.31	0.79	1.0	0.45	0.26	0.32	0.48
25	0.88	2.6	3.51	1.87	1.8	2.5	0.7	0.82	1.07
36	0.48	1.06	1.52	0.77	0.66	0.89	0.28	0.3	0.48
52	1.0	1.95	2.9	1.11	1.1	1.02	0.36	0.48	0.54
72	5.56	7.32	9.4	2.16	1.88	1.9	0.9	1.1	1.26
100	62.15	30.05	29.32	14.4	12.04	13.06	9.6	9.62	10.11
150	27.71	45.6	41.62	55.26	56.12	56.0	62.02	63.0	59.3
200	1.09	5.1	4.69	13.35	16.5	12.41	20.0	18.02	20.65
240	0.2	0.41	0.4	2.89	2.89	3.0	3.33	3.2	3.41

Sieve Size	Percentage								
	Sample Number								
	W64	W65	W66	W67	W68	W69	W70	W71	W72
8	0.03	0.06	0.02	0.37	0.15	0.08	0.09	0.1	0.5
14	0.45	0.6	0.28	0.6	0.41	0.42	0.69	0.68	0.22
25	1.39	2.31	1.63	1.6	1.37	1.25	2.38	1.29	2.12
36	0.9	1.3	1.2	1.0	0.89	0.99	0.9	0.55	0.72
52	3.63	4.43	3.34	2.9	2.4	2.72	1.78	1.67	1.71
72	16.52	15.21	15.01	11.0	9.4	10.3	9.02	10.99	9.48
100	22.81	20.2	21.77	18.58	17.15	18.3	23.75	26.29	24.6
150	27.43	27.6	27.93	48.7	50.45	49.98	32.5	31.7	31.91
200	11.02	9.67	9.7	9.89	11.43	10.78	10.22	10.81	12.0
240	4.05	4.91	4.5	2.41	2.86	2.46	4.63	4.63	5.05

Sieve Size	Percentage								
British Standard	Sample Number								
	W73	W74	W75	W76	W77	W78	W79	W80	W81
8	0.15	0.02	0.1	0.01	0.11	-	-	0.15	0.2
14	0.23	0.2	0.75	1.1	0.37	0.28	0.5	0.6	0.5
25	1.92	1.88	2.0	4.22	2.2	2.91	2.02	1.65	1.85
36	0.7	0.78	0.75	1.2	0.75	1.1	0.72	0.61	1.98
52	1.6	1.7	1.91	2.09	1.55	2.11	1.7	1.6	5.95
72	8.85	9.35	10.74	8.97	8.41	10.19	10.4	9.18	18.91
100	23.35	23.52	25.05	22.44	22.25	24.15	25.86	24.5	3.04
150	33.3	30.9	32.08	28.32	33.25	30.2	31.1	33.8	34.7
200	11.25	11.48	11.21	10.84	11.15	8.02	9.9	10.81	2.6
240	5.5	5.32	4.22	4.95	5.19	5.78	4.5	5.1	0.58

Sieve Size	Percentage								
	Sample Number								
	W82	W83	W84	W85	W86	W87	W88	W89	W90
8	0.07	0.08	0.1	0.1	0.09	0.04	0.02	0.17	0.4
14	0.3	0.3	0.42	0.39	0.3	0.51	0.12	0.52	0.88
25	1.22	1.73	1.89	1.82	1.7	1.92	1.0	3.1	4.47
36	1.84	2.88	1.92	4.38	4.45	4.09	3.55	2.1	3.28
52	6.35	8.21	6.34	19.0	18.9	18.89	17.81	3.55	5.4
72	21.03	24.26	19.48	44.32	45.3	45.81	43.62	5.12	6.68
100	32.34	30.0	31.79	23.51	23.48	22.82	26.89	9.9	10.42
150	32.13	28.07	33.78	5.01	4.93	4.8	5.95	37.0	27.78
200	2.1	2.08	2.51	0.03	0.1	0.09	0.1	19.2	16.18
240	0.62	0.41	0.3	0.2	0.12	0.2	0.27	4.66	4.1

Sieve Size	Percentage								
British Standard	Sample Number								
	W91	W92	W93	W94	W95	W96	W97	W98	W99
8	0.39	0.52	0.18	0.11	0.01	0.01	0.21	0.1	0.17
14	0.58	1.15	0.9	0.89	0.6	0.48	2.6	0.87	0.71
25	2.9	2.81	2.3	2.51	2.64	2.4	4.65	2.31	1.49
36	1.98	1.29	1.0	0.9	1.21	0.95	1.32	0.89	0.7
52	2.6	1.95	1.38	1.26	1.53	1.35	1.6	1.1	1.04
72	3.36	3.2	2.65	2.39	2.71	2.72	3.55	2.78	2.5
100	12.0	13.95	12.88	14.01	13.5	13.32	26.92	13.75	14.7
150	39.75	41.09	42.41	41.49	41.53	41.83	40.62	44.6	44.32
200	17.48	16.51	16.78	16.39	19.1	17.52	9.55	15.15	16.87
240	3.61	2.71	3.63	3.7	2.83	2.78	0.6	2.8	2.98

Sieve Size	Percentage								
	Sample Number								
	W100	W101	W102	W103	W104	W105	W106	W107	W108
8	-	-	0.02	-	0.09	0.08	0.02	0.13	0.21
14	0.55	0.22	0.38	0.32	0.11	0.37	0.27	0.3	0.49
25	2.41	1.39	1.49	0.92	1.92	1.25	0.9	0.81	0.9
36	1.35	0.87	0.77	1.2	2.3	2.2	2.01	0.52	0.52
52	1.25	0.88	0.7	1.41	15.89	16.12	14.41	1.33	1.32
72	3.13	2.92	2.42	3.45	50.6	49.92	47.78	16.14	15.49
100	15.26	15.96	16.6	16.32	23.7	24.4	27.33	32.12	31.29
150	44.55	44.48	45.61	41.7	3.3	4.6	6.28	42.27	43.1
200	13.82	16.6	14.59	15.52	0.15	0.23	0.32	4.2	4.32
240	3.37	3.0	3.49	2.38	0.2	0.02	0.2	0.46	0.47

Sieve Size	Percentage								
British Standard	Sample Number								
	W109	W110	W111	W112	W113	W114	W115	W116	W117
8	0.07	0.26	0.16	0.22	1.01	0.1	0.21	0.24	-
14	0.29	0.49	0.29	0.5	0.61	0.13	0.69	0.41	0.3
25	0.77	1.0	0.7	1.1	1.29	0.46	1.8	1.11	1.66
36	0.43	0.52	0.39	0.59	0.5	0.32	1.0	1.0	1.0
52	1.3	1.4	0.63	0.89	0.59	0.47	1.2	1.42	1.3
72	17.53	17.48	9.41	11.58	8.23	7.75	13.42	10.59	12.1
100	32.88	33.13	67.56	63.93	61.91	66.5	58.86	62.9	58.8
150	40.61	39.67	19.71	19.8	24.11	22.89	20.9	20.66	22.3
200	3.7	3.7	0.5	0.55	0.62	0.62	1.0	0.91	1.31
240	0.46	0.31	0.05	0.1	0.1	0.1	0.13	0.15	0.2

Sieve Size		Sample Number								
		W118	W119	W120	W121	W122	W123	W124	W125	W126
		W118	W119	W120	W121	W122	W123	W124	W125	W126
8		0.25	0.13	0.41	0.4	0.55	9.67	32.72	6.7	4.4
14		0.4	0.9	1.4	0.8	1.92	14.2	13.28	12.82	7.12
25		1.35	5.67	9.46	6.83	10.88	18.26	15.85	29.19	22.12
36		0.87	10.1	14.39	12.32	15.02	10.95	5.91	16.6	17.11
52		1.03	21.89	23.0	25.23	23.89	12.25	5.27	13.71	16.05
72		11.97	38.91	33.75	38.59	34.32	16.68	6.5	12.5	18.6
100		59.1	17.23	13.5	12.83	10.8	8.04	6.28	5.37	9.7
150		22.41	3.63	2.77	2.06	1.65	4.42	5.1	1.92	3.36
200		1.26	0.2	0.12	0.1	0.08	3.11	2.9	0.18	0.27
240		0.18	0.01	0.01	0.01	0.01	0.6	0.5	0.02	0.05

Sieve Size	Percentage								
British Standard	Sample Number								
	W127	W128	W129	W130	W131	W132	W133	W134	W135
8	0.2	0.17	0.02	-	-	-	0.59	0.01	0.5
14	0.67	0.49	0.2	0.26	0.13	0.16	0.32	0.32	0.51
25	1.21	2.71	0.6	0.97	0.7	1.1	1.83	1.5	1.28
36	2.4	7.8	0.4	0.58	0.33	0.65	1.1	0.97	0.81
52	17.57	22.55	0.69	0.88	0.53	1.01	1.9	1.6	1.4
72	52.2	47.2	16.6	17.3	15.4	17.8	14.3	12.32	12.3
100	22.55	15.11	64.99	63.09	64.76	63.95	53.58	54.6	54.9
150	2.62	2.1	15.58	15.75	16.7	14.02	22.9	25.11	24.5
200	0.11	0.17	0.47	0.48	0.6	0.42	1.96	2.17	2.4
240	0.01	0.08	0.03	0.03	0.06	0.03	0.22	0.25	0.3

Sieve Size	Percentage				
	Sample Number				
	W136	W137	W138	W139	W140
8	0.01	0.12	-	-	-
14	0.27	0.12	0.2	0.1	0.09
25	1.08	0.65	0.9	0.5	0.5
36	0.7	0.38	0.4	0.31	0.31
52	1.25	0.4	0.4	0.32	0.39
72	11.45	1.8	2.1	1.76	2.13
100	53.4	30.47	32.98	32.53	33.42
150	28.0	54.4	53.17	54.56	53.62
200	2.53	9.49	8.1	8.1	7.71
240	0.31	0.72	0.59	0.55	0.59

Sieve Size	Percentage								
British Standard	Sample Number								
	D1	D2	D3	D4	D5	D6	D7	D8	D9
8	-	-	-	-	-	-	-	-	-
14	-	-	-	-	-	-	-	0.02	-
25	0.2	-	0.1	0.1	0.04	0.1	0.1	0.12	0.1
36	0.2	0.1	0.15	0.2	0.2	0.8	1.04	1.0	1.0
52	7.0	4.24	5.62	10.8	8.24	36.24	37.0	26.4	25.8
72	61.56	54.8	58.18	62.44	61.1	57.2	56.8	61.96	62.0
100	27.54	36.6	32.07	24.4	27.66	4.58	4.4	9.6	10.2
150	1.76	3.0	2.38	1.4	1.8	0.2	0.2	0.20	0.3
200	0.02	0.2	0.11	-	-	-	-	-	-
240	-	-	-	-	-	-	-	-	-

Sieve Size	Percentage			
	Sample Number			
	D10	D11	D12	D13
8	-	-	-	-
14	-	-	-	-
25	-	-	0.02	0.04
36	0.1	0.1	0.42	0.5
52	9.24	8.52	10.64	10.4
72	58.44	59.7	59.76	60.9
100	29.04	29.3	26.38	25.5
150	1.84	1.84	1.90	2.2
200	0.1	0.08	0.12	0.18
240	0.06	0.06	0.04	0.06

[illegible][illegible]

Sieve Size British Standard	Percentage					
	Sample Number					
	B37	B38	B39	B40	B41	B42
8	0.08	0.18	0.36	0.18	0.2	0.1
14	0.18	0.18	0.36	0.2	0.1	0.16
25	0.66	0.62	1.36	1.14	0.64	1.0
36	1.26	1.22	1.64	1.58	0.9	1.22
52	10.34	10.44	9.14	8.62	6.4	7.44
72	37.22	36.8	46.28	43.4	34.8	38.96
100	38.1	39.02	33.44	36.58	43.72	40.56
150	11.52	11.30	7.18	7.86	12.84	10.2
200	0.14	0.14	0.18	0.12	0.16	0.1
240	0.04	0.10	0.06	0.04	0.02	0.02

Sieve Size	Percentage								
British Standard	Sample Number								
	R1	R2	R3	R4	R5	R6	R7	R8	R9
8	0.86	1.0	0.1	0.04	3.76	1.41	2.7	1.63	0.8
14	1.82	2.98	0.3	0.2	6.1	3.01	3.8	2.96	3.16
25	27.54	33.1	5.9	6.04	27.0	24.78	21.3	19.92	31.18
36	29.6	28.96	10.38	11.62	17.6	21.8	17.98	19.32	25.68
52	23.4	20.9	22.36	24.6	13.96	20.78	15.68	20.93	18.14
72	9.2	7.2	23.62	23.56	7.6	11.27	9.04	12.84	6.6
100	3.24	3.22	16.24	15.4	4.56	6.11	5.84	6.74	3.06
150	1.76	1.41	10.78	10.12	4.0	4.62	4.84	6.06	2.72
200	0.46	0.36	3.70	3.0	1.56	1.77	1.94	2.39	1.16
240	0.22	0.2	1.62	1.3	0.82	0.92	1.04	1.21	0.68

Sieve Size	Percentage		
	Sample Number		
	R10	R11	R12
8	2.21	0.34	0.56
14	5.23	0.76	1.39
25	39.11	14.4	19.37
36	24.58	23.04	26.13
52	14.89	27.86	27.28
72	4.89	14.62	12.2
100	2.23	6.81	5.04
150	3.9	4.97	3.24
200	0.8	1.92	1.11
240	0.42	1.0	0.58

Sieve Size	Percentage								
British Standard	Sample Number								
	U1	U2	U3	U4	U5	U6	U7	U8	U9
8	0.4	0.6	0.4	1.86	0.4	0.12	-	-	-
14	0.84	1.0	0.84	0.84	0.24	0.18	-	-	0.02
25	10.0	9.3	13.6	12.84	4.26	4.0	0.04	0.04	0.1
36	24.2	21.8	16.5	16.9	10.9	10.4	0.06	0.06	0.1
52	37.4	36.3	37.7	38.42	48.2	48.74	0.22	0.22	0.38
72	17.30	18.84	21.26	21.06	24.4	25.16	2.7	2.8	4.2
100	5.04	6.26	4.4	4.0	5.52	5.56	16.2	17.5	40.2
150	1.7	2.4	1.74	1.4	2.44	2.4	35.02	34.0	44.64
200	0.2	0.4	0.4	0.2	0.6	0.6	16.16	16.0	6.84
240	-	-	0.12	-	0.1	0.1	0.2	3.22	1.4

Sieve Size	Percentage								
	Sample Number								
	U10	U11	U12	U13	U14	U15	U16	U17	U18
8	-	-	0.18	0.2	-	3.4	2.8	0.92	0.76
14	0.04	0.12	0.16	0.24	0.1	5.64	4.82	1.34	1.44
25	0.04	0.22	0.4	0.3	0.36	7.0	6.94	19.84	22.72
36	0.1	0.22	0.38	0.32	0.4	4.22	4.1	34.54	35.9
52	0.32	1.8	2.22	3.0	3.44	10.0	9.76	28.0	26.44
72	2.8	13.12	15.0	20.0	21.0	19.04	19.14	8.36	7.56
100	31.78	31.8	31.34	33.8	32.72	20.16	20.44	2.28	2.4
150	51.62	35.96	35.34	24.8	25.4	17.8	18.56	1.36	1.16
200	8.2	9.4	9.0	7.32	6.78	5.62	6.0	0.42	0.4
240	2.42	3.16	2.8	3.2	3.0	2.24	2.24	0.28	0.16

Sieve Size	Percentage								
British Standard	Sample Number								
	U19	U20	U21	U22	U23	U24	U25	U26	U27
8	0.6	0.64	0.2	0.24	-	-	2.54	2.6	1.58
14	0.9	1.0	0.38	0.4	0.12	0.18	2.2	1.6	2.2
25	8.9	10.22	5.56	6.9	3.92	4.42	10.44	8.62	6.4
36	20.64	22.38	21.5	24.2	23.6	25.78	15.02	13.76	10.98
52	39.76	40.1	42.0	43.34	48.24	48.68	26.8	26.16	26.76
72	17.5	16.18	18.24	15.84	15.84	14.6	21.32	22.0	23.46
100	5.6	4.82	6.42	4.8	3.84	3.42	11.24	12.5	13.6
150	2.46	1.94	2.8	1.72	1.6	1.2	5.48	6.6	8.04
200	0.72	0.54	0.78	0.4	0.5	0.42	1.42	1.84	2.3
240	0.28	0.26	0.22	0.24	0.34	0.14	0.6	0.76	0.96

Sieve Size	Percentage								
	Sample Number								
	U28	U29	U30	U31	U32	U33	U34	U35	U36
8	1.4	2.0	1.3	0.08	0.1	0.18	0.06	-	-
14	1.5	1.96	2.26	0.14	0.06	0.2	0.08	-	-
25	5.48	9.76	10.16	0.26	0.33	0.42	0.36	0.24	0.1
36	11.22	13.62	13.64	0.32	1.16	0.36	1.0	0.32	0.2
52	26.28	25.2	24.4	3.1	8.44	3.5	7.86	2.0	2.02
72	23.16	21.0	21.18	19.4	24.6	20.58	24.36	14.0	15.5
100	14.02	12.76	13.30	33.14	30.44	34.64	30.56	25.56	28.2
150	8.46	7.5	7.6	27.46	22.64	27.78	23.46	28.36	31.38
200	2.64	2.18	2.34	7.36	6.64	7.6	7.2	10.9	12.62
240	1.18	1.0	0.96	2.78	2.8	3.08	2.98	5.16	5.54

Sieve Size	Percentage								
British Standard	Sample Number								
	U37	U38	U39	U40	U41	U42	U43	U44	U45
8	0.2	0.06	0.6	0.2	0.2	0.02	0.4	0.26	0.04
14	0.2	0.2	0.46	0.6	0.24	0.06	0.2	0.44	0.16
25	0.48	0.68	2.24	0.6	0.78	0.84	1.22	1.64	0.66
36	1.84	1.52	6.0	6.22	3.06	3.44	4.42	5.0	3.02
52	9.62	11.26	21.8	22.24	18.18	19.97	23.5	25.0	19.8
72	35.74	35.1	40.36	39.9	43.24	44.2	41.3	40.56	44.88
100	28.96	28.6	19.04	19.36	23.24	20.6	18.56	18.2	20.84
150	13.78	13.3	5.6	5.56	7.36	6.64	6.0	5.42	6.56
200	3.8	3.6	1.14	1.04	1.5	1.4	1.4	1.14	1.36
240	1.4	1.2	0.44	0.4	0.2	0.2	0.6	0.38	0.42

Sieve Size	Percentage								
	Sample Number								
	U46	U47	U48	U49	U50	U51	U52	U53	U54
8	0.24	0.14	0.6	-	-	-	0.12	-	-
14	0.16	0.2	0.24	0.1	0.06	-	0.06	-	-
25	0.82	0.8	0.98	0.12	0.12	0.02	0.1	0.02	0.1
36	3.4	2.94	3.4	0.36	0.42	0.3	0.4	0.2	0.2
52	21.16	18.92	19.44	14.36	14.36	12.18	14.26	8.6	10.4
72	44.02	44.16	44.48	53.02	51.4	51.82	54.06	51.5	54.02
100	19.4	23.0	22.0	25.7	26.42	27.78	25.7	31.5	29.3
150	6.44	6.5	6.16	4.8	5.6	6.02	4.56	6.7	5.18
200	1.54	1.12	1.78	0.4	0.68	0.76	0.4	0.8	0.42
240	0.6	0.4	0.2	0.2	0.28	0.28	0.2	0.34	0.16

Sieve Size British Standard	Percentage								
	Sample Number								
	U55	U56	U57	U58	U59	U60	U61	U62	U63
8	-	-	-	-	-	-	-	-	-
14	-	-	-	-	-	0.04	-	-	-
25	-	-	0.02	-	0.04	0.04	-	0.04	0.16
36	0.1	0.06	0.16	0.16	0.38	0.32	0.02	0.1	0.24
52	5.2	5.58	7.16	9.0	13.4	11.96	0.14	0.26	0.4
72	47.4	48.22	50.6	53.4	52.84	53.58	0.4	0.36	0.42
100	35.5	35.2	32.44	29.48	25.78	26.8	12.62	12.56	18.1
150	9.04	8.58	7.4	5.98	5.44	5.42	45.4	46.04	47.74
200	1.5	1.2	1.0	0.64	0.7	0.6	22.26	21.72	18.7
240	0.6	0.4	0.4	0.26	0.26	0.22	13.16	12.8	10.2

Sieve Size	Sample Number								
	U64	U65	U66	U67	U68	U69	U70	U71	U72
	U64	U65	U66	U67	U68	U69	U70	U71	U72
8	-	-	-	-	-	-	-	-	-
14	0.14	-	-	-	-	-	-	-	-
25	0.34	0.1	0.06	0.02	0.04	0.04	0.01	0.04	0.04
36	0.36	0.1	0.1	0.06	0.06	0.06	0.02	0.04	0.04
52	0.36	0.2	0.2	0.1	0.14	0.1	0.02	0.04	0.04
72	0.6	0.42	0.36	0.36	0.42	0.2	0.2	0.3	0.2
100	14.6	12.2	10.44	13.58	12.24	8.2	9.1	9.8	10.7
150	45.42	47.2	43.8	47.42	48.6	42.8	41.9	42.06	41.2
200	21.24	21.6	22.16	23.24	23.16	23.0	22.6	21.36	20.42
240	11.9	13.38	13.84	12.78	12.78	13.84	13.78	12.38	12.04

Sieve Size British Standard	Percentage								
	Sample Number								
	U73	U74	U75	U76	U77	U78	U79	U80	U81
8	0.3	0.22	0.4	0.06	0.42	0.28	0.3	0.3	1.08
14	9.72	7.04	8.8	4.16	11.6	5.6	1.42	0.66	1.62
25	28.06	23.1	24.82	16.54	30.0	21.44	19.7	14.52	14.9
36	22.64	22.38	19.4	17.56	20.16	19.1	13.7	15.9	14.3
52	24.56	26.72	18.72	20.56	17.38	20.6	13.49	17.01	16.6
72	9.9	12.44	12.58	16.62	9.8	15.0	11.1	12.63	13.1
100	2.58	4.04	7.6	11.64	5.24	9.32	9.45	9.52	10.5
150	0.86	2.0	4.5	7.4	1.82	5.98	12.48	11.77	13.4
200	0.2	0.52	0.24	2.4	1.04	1.4	5.2	7.49	5.39
240	0.12	0.24	1.70	1.2	0.14	0.8	4.9	1.52	1.62

Sieve Size	Percentage							
	Sample Number							
	U82	U83	U84	U85	U86	U87	U88	U89
8	8.08	0.4	0.5	1.1	-	-	11.41	10.1
14	5.12	1.4	1.0	1.51	-	-	2.2	3.4
25	14.28	18.6	14.38	15.1	0.23	0.1	6.4	7.1
36	7.8	14.2	15.4	13.9	0.20	0.31	13.1	12.8
52	16.2	14.0	18.1	16.82	2.9	2.32	16.9	15.78
72	29.3	11.4	12.21	13.3	16.4	15.5	23.2	20.3
100	13.3	9.9	9.41	10.1	28.2	28.7	13.84	16.9
150	3.2	11.8	11.88	13.5	26.4	32.0	6.32	8.5
200	0.42	4.9	7.3	4.0	13.8	12.6	2.1	3.61
240	0.18	4.8	1.6	1.1	6.4	5.7	0.9	1.4

Samples Used for Factor Analysis 2

<u>Seaton Carew Beach 2</u>	<u>Saltburn Beach</u>	<u>Beach N.of Saltburn</u>	<u>Marske Beach 1</u>	<u>South Gare Beach 10</u>	<u>Redcar Beach 3</u>
-	0.39	-	0.16	-	0.64
-	0.3	0.28	0.02	-	0.8
0.01	2.3	1.07	0.76	0.06	1.4
0.02	6.11	4.52	0.98	0.12	5.41
0.2	14.6	18.32	4.78	3.8	1.7
11.2	36.33	43.29	34.36	35.2	13.82
62.55	37.09	30.8	54.36	54.81	65.78
25.2	2.3	1.49	4.36	5.91	14.78
0.5	0.03	-	0.04	0.07	0.2
0.1	-	-	-	-	-

<u>Musselborough Beach 1</u>	<u>Hartlepool Yacht Club 3</u>	<u>Hartlepool Yacht Club 1</u>	<u>Crimdon Beach 1</u>	<u>Crimdon Beach 3</u>
0.32	0.1	0.18	0.36	0.18
1.4	0.16	0.18	0.36	0.6
4.09	1.0	0.62	2.64	5.58
1.22	1.22	17.66	14.72	10.34
10.66	7.44	10.44	52.26	39.24
33.22	38.96	36.8	23.0	33.4
41.52	40.56	39.02	2.0	5.22
2.32	10.2	11.3	0.04	0.3
0.01	0.1	0.14	-	-
-	0.02	0.1	-	-

<u>Droman Beach A</u>	<u>Droman Beach B</u>	<u>S. Gare Dune D</u>	<u>S. Gare Dune A</u>	<u>Cliffslip Huntcliff</u>	<u>Musselborough Sand Dune</u>
-	-	-	-	-	0.1
0.27	1.4	-	-	0.07	0.17
2.9	13.59	0.02	0.01	0.45	2.46
10.34	19.12	0.28	0.11	1.44	3.99
23.58	29.01	5.21	2.63	8.42	7.09
44.18	27.62	48.04	35.02	38.8	28.73
5.22	17.05	8.54	43.68	58.51	46.85
0.3	1.39	0.51	2.45	3.5	2.72
-	0.01	0.01	0.04	0.02	0.08
-	-	0.01	-	-	0.01

<u>Crimdon Dune 2</u>	<u>Crimdon Dune 18'</u>	<u>Crimdon Dune 33'</u>	<u>Crimdon Dune 5'</u>	<u>Droman Dune B</u>
-	-	-	-	-
0.02	-	-	-	0.07
0.12	0.05	0.1	0.2	1.5
1.0	0.8	0.2	0.2	8.35
26.4	36.24	10.8	7.0	22.98
61.96	57.2	62.44	61.56	43.8
53.87	9.6	4.58	24.4	27.54
3.1	0.26	0.2	1.4	1.76
0.07	-	-	-	0.02
-	-	-	-	-

<u>Droman</u> <u>Dune A</u>	<u>Saltburn</u> <u>River Sand</u>	<u>Thorpe</u> <u>Thewles 4</u>	<u>Saltburn</u> <u>1967 Flood</u>	<u>Thorpe</u> <u>Thewles 1</u>	<u>Thorpe</u> <u>Thewles 2</u>
-	-	-	4.07	1.41	1.63
0.01	0.06	0.05	6.61	3.01	2.96
1.55	1.01	0.32	29.27	24.78	19.92
9.53	5.42	5.18	19.08	21.8	19.32
24.85	18.92	18.3	15.13	20.78	20.93
42.08	35.51	36.38	8.24	11.27	12.84
18.98	24.03	24.37	4.94	6.11	6.74
2.69	10.42	10.12	4.33	4.62	6.06
0.27	2.49	2.4	1.69	1.77	2.39
-	0.83	0.93	0.89	1.15	1.21

<u>Billingham</u> <u>Beck 1</u>	<u>Billingham</u> <u>Beck 2</u>	<u>Offshore</u> <u>SLS 10</u>	<u>Offshore</u> <u>SLS 4</u>	<u>Offshore</u> <u>LOC 2</u>
0.86	0.04	0.01	0.05	0.08
1.82	0.2	0.02	0.29	0.08
27.54	6.04	0.54	1.28	1.21
29.6	11.62	0.38	1.59	0.4
23.4	24.6	0.61	5.9	0.78
9.2	23.56	1.8	18.93	2.8
3.24	15.4	14.53	19.2	23.42
1.76	10.12	48.63	34.64	53.57
0.46	3.0	19.05	11.26	8.95
0.22	1.3	6.54	3.08	2.26

<u>Offshore</u> <u>SLS 3</u>	<u>Offshore</u> <u>LOC 12</u>	<u>Offshore</u> <u>R5</u>	<u>Offshore</u> <u>R1</u>	<u>Offshore</u> <u>LOC 16</u>	<u>Offshore</u> <u>SLS 9</u>	<u>Offshore</u> <u>R2</u>
0.06	0.2	2.3	1.01	0.02	0.27	0.25
0.6	0.45	1.3	0.61	0.26	0.52	0.04
2.31	2.6	1.13	1.29	2.26	1.3	1.35
1.3	2.89	0.8	0.5	4.89	0.48	0.87
4.43	13.49	1.86	0.59	18.29	0.7	1.03
15.21	46.15	15.32	8.23	41.85	2.8	11.97
2.02	28.91	59.59	61.91	27.23	18.61	59.1
27.6	4.09	15.9	24.11	4.33	55.79	22.41
9.67	0.1	0.58	0.62	0.11	15.3	1.26
4.91	0.01	0.09	0.1	0.01	2.09	0.18

<u>Offshore</u> <u>R8</u>	<u>Offshore</u> <u>LOC 7</u>	<u>Offshore</u> <u>LOC 1</u>	<u>Offshore</u> <u>SLS 6</u>	<u>Offshore</u> <u>LOC 10</u>	<u>Offshore</u> <u>R9</u>	<u>Droman</u> <u>Offshore</u> <u>A Site 1</u>
-	0.19	0.06	0.08	0.09	-	-
0.09	0.47	0.39	0.42	0.3	0.26	3.2
0.5	1.95	1.95	1.25	1.7	0.97	9.09
0.31	2.25	1.12	0.99	4.45	0.58	13.8
0.39	7.45	2.48	2.72	18.9	0.88	27.2
2.13	23.9	6.2	10.3	45.3	17.3	34.33
33.42	37.13	8.0	18.3	23.48	63.09	10.58
53.62	23.61	31.07	49.98	4.93	15.75	1.5
7.71	1.58	18.5	10.78	0.1	0.48	0.25
0.59	0.11	10.39	2.46	0.12	0.03	0.05

<u>Droman Offshore B Site 1</u>	<u>Droman Offshore A Site 2</u>	<u>Droman Offshore B Site 2</u>	<u>Droman Offshore C Site 2</u>	<u>Coatham Stob Yellow Sand 1</u>	<u>Coatham Stob Yellow Sand 3</u>
-	0.01	-	-	0.1	0.01
1.9	1.01	0.83	0.68	0.06	0.02
6.51	3.65	5.01	4.7	0.41	0.7
11.65	9.59	12.4	12.41	2.71	3.95
25.28	24.2	27.71	25.81	8.92	12.22
36.3	40.93	37.28	37.88	26.5	29.65
15.55	17.6	13.3	14.7	31.97	28.2
2.21	2.75	2.8	3.15	18.71	15.65
0.31	0.36	0.51	0.42	5.34	4.29
0.1	0.1	0.11	0.09	1.51	1.42

<u>Coatham Stob 1</u>	<u>Coatham Stob 4</u>	<u>Coatham Stob 4 2</u>	<u>Coatham Stob 4 1</u>	<u>Coatham Stob 5 2</u>	<u>Coatham Stob Same as 3</u>	<u>Coatham Stob 3</u>
11.53	2.5	0.12	0.3	12.5	0.67	-
2.22	0.78	0.42	0.61	2.7	0.5	0.07
6.24	4.9	1.37	1.38	10.53	0.78	0.04
13.12	8.78	2.8	2.0	23.2	4.4	0.85
16.88	12.95	9.36	5.78	23.37	17.58	4.48
23.2	21.3	31.19	20.72	17.8	39.38	22.09
13.71	21.6	32.07	35.16	6.61	23.66	33.85
6.4	15.79	15.57	22.4	1.81	8.6	22.95
2.08	4.8	3.6	5.8	0.4	1.83	7.12
0.82	1.6	1.03	1.88	0.15	0.58	2.31

<u>Brierton Top 3A</u>	<u>Brierton 2C</u>	<u>Brierton Mid 3B</u>	<u>Brierton 3C Bottom</u>	<u>Brierton C4</u>
1.92	0.28	0.04	0.02	0.04
0.76	5.6	0.08	0.02	0.16
1.2	21.44	0.4	0.08	0.66
2.6	19.10	0.8	0.18	3.02
18.16	20.6	6.22	1.8	19.8
32.84	15.0	19.16	15.0	44.88
23.64	9.32	21.84	26.2	20.84
11.78	5.98	22.76	27.0	6.56
2.8	1.4	10.02	11.2	1.36
1.2	0.8	5.94	6.18	0.42

<u>Brierton 4C Above 3C</u>	<u>Brierton A2</u>	<u>Brierton C2</u>	<u>Brierton C5</u>
0.18	1.58	0.6	0.14
0.2	2.2	0.46	0.2
1.22	6.4	2.24	0.8
2.42	10.98	6.0	2.94
12.94	26.76	21.8	18.92
31.56	23.46	40.36	44.16
28.92	13.6	19.04	23.0
15.2	8.04	5.6	6.5
3.3	2.3	1.14	1.12
1.2	0.96	0.44	0.4

<u>Wynyard Basal Sand</u>	<u>Wynyard C</u>	<u>Wynyard A</u>	<u>Wynyard 3</u>	<u>Wynyard 2</u>
-	0.4	0.4	0.2	0.6
-	0.84	0.84	0.38	0.9
0.04	10.0	13.6	5.56	8.9
0.06	24.2	16.5	21.5	20.64
0.22	37.4	37.7	42.0	39.76
2.7	17.3	21.26	18.24	17.5
16.2	5.04	4.4	6.42	5.6
35.02	1.7	17.4	2.8	2.46
16.16	0.2	0.4	0.78	0.72
0.2	-	0.12	0.22	0.28

<u>Hurworth 3</u>	<u>Hurworth H3</u>	<u>Hurworth 1</u>	<u>Hurworth 5</u>	<u>Hurworth H6</u>
-	-	-	-	-
-	-	-	-	-
0.1	0.02	-	0.04	0.04
0.2	0.2	0.02	0.07	0.38
0.24	8.6	0.14	0.11	13.4
0.44	51.5	0.42	0.22	52.84
13.1	31.5	13.38	9.16	25.78
49.47	6.7	48.12	47.83	5.44
22.61	0.8	23.59	25.7	0.7
14.02	0.34	13.95	15.47	0.26

<u>Newton Bewley 2</u>	<u>Girsby Flat</u>	<u>Eaglescliffe 3</u>	<u>Eaglescliffe 4</u>	<u>Rockcliffe 1</u>	<u>Stockton Norton 1</u>
0.3	-	0.05	0.02	0.2	-
0.66	1.36	0.1	0.02	0.22	0.07
14.52	5.4	0.2	0.1	0.31	15.4
15.9	5.2	0.2	0.11	0.88	38.8
17.01	7.2	0.41	0.22	3.0	20.68
12.63	15.53	1.85	1.88	27.01	12.38
9.52	19.33	9.8	23.79	46.62	4.55
11.77	17.78	44.32	56.35	13.63	2.09
7.49	8.3	23.22	12.6	3.25	1.03
1.52	3.28	7.22	2.1	1.01	0.51

<u>Ireland Interglac Beach</u>	<u>Edinburgh Raised Sand</u>	<u>Edinburgh Fluvio- Glacial</u>	<u>Edinburgh Raised Marine</u>	<u>Edinburgh Raised Marine</u>
-	-	-	-	-
0.05	0.01	-	0.01	-
0.96	0.02	0.1	0.06	0.18
2.32	0.11	0.24	0.06	0.22
8.5	0.92	1.28	0.07	0.23
21.31	16.62	15.77	1.6	4.08
24.26	37.31	38.8	23.7	26.91
23.8	27.49	31.07	45.6	48.92
10.3	11.29	7.8	21.95	13.31
-	2.1	2.14	3.27	2.51

APPENDIX 5Mineralogical DataSamples used in Factor Analysis 2

1) <u>Heavy Minerals</u>		(Weights in Grams)			
<u>Beach</u>	<u>Weight</u>	<u>Weight</u> <u>After</u> <u>Hand</u> <u>Magnet</u>	<u>Weight</u> <u>After</u> <u>Electro-</u> <u>Magnet</u>	<u>%</u> <u>Weight</u> <u>Loss</u>	
<u>Seaton Carew Beach 2</u>					
<u>B. S. Sieves</u>					
72	22.859	22.843	22.643	0.945	
100	61.53	61.477	60.640	1.419	
150	12.674	12.612	11.742	7.354	
200	--	--	--	--	
<u>Saltburn Beach</u>					
72	36.317	36.271	35.427	2.451	
100	46.561	46.550	45.513	2.251	
150	2.453	2.411	2.025	17.448	
200	--	--	--	--	
<u>Beach North of Saltburn</u>					
72	41.501	41.475	40.740	1.834	
100	21.570	21.560	21.064	2.346	
150	0.603	0.600	0.545	9.618	
200	0.029	0.028	0.021	27.586	
<u>Marske Beach (1)</u>					
72	32.300	32.265	31.724	1.783	
100	34.842	34.807	34.379	1.329	
150	2.016	2.008	1.849	8.284	
200	--	--	--	--	
<u>South Gare 10</u>					
72	38.506	38.490	38.081	1.104	
100	52.395	52.379	51.889	0.966	
150	4.654	4.650	4.226	9.196	
200	--	--	--	--	

	Weight	Weight After Hand Magnet	Weight After Electro- Magnet	% Weight Loss
<u>Redcar Beach (3)</u>				
72	26.014	26.007	25.697	1.218
100	56.864	56.830	56.308	0.978
150	7.819	7.812	6.974	10.807
200	--	--	--	--
<u>Musselborough Beach</u>				
72	38.732	38.621	36.694	5.262
100	42.550	42.464	41.166	3.253
150	2.460	2.397	2.060	16.260
<u>H.Y.C. Beach 3</u>				
72	41.114	40.990	40.219	2.177
100	39.735	39.614	38.669	2.683
150	10.388	10.273	9.914	4.563
<u>H.Y.C. Beach 1</u>				
72	40.507	40.352	39.618	2.195
100	37.691	37.623	36.760	2.470
150	10.106	10.076	9.575	5.254
<u>Crimdon Beach 1</u>				
72	28.432	28.399	27.807	2.198
100	3.641	3.630	3.463	4.889
150	0.076	0.076	0.054	28.947
<u>Crimdon Beach 3</u>				
72	35.265	35.240	34.536	2.067
100	5.027	5.017	4.739	5.729
150	0.175	0.172	0.136	22.286
<u>Droman Beach A</u>				
72	44.168	44.167	44.045	8.278
100	17.067	17.067	16.978	0.521
150	1.379	1.379	1.337	3.045
200	0.781	0.781	0.467	41.844

	<u>Weight</u>	<u>Weight After Hand Magnet</u>	<u>Weight After Electro- Magnet</u>	<u>% Weight Loss</u>
S. Gare Dune D				
72	42.570	42.550	42.209	0.848
100	50.477	50.453	50.085	0.776
150	2.249	2.238	2.166	3.690
S. Gare Dune A				
72	51.254	51.210	50.527	1.418
100	39.725	39.685	39.221	1.269
150	2.039	2.017	1.945	4.610
Cliff Slip. Huntcliff				
72	40.100	40.089	39.503	1.489
100	50.448	50.437	49.755	1.374
150	2.895	2.870	2.541	12.228
Musselborough Sand Dune				
72	40.379	40.292	39.081	3.214
100	44.123	44.002	42.501	3.676
150	1.704	1.634	1.316	22.770
200	0.107	0.092	0.052	51.402
Crimdon Dune 2				
72	64.030	64.003	63.376	1.021
100	24.977	24.951	24.228	2.999
150	1.275	1.253	1.098	13.882
Crimdon Dune 18'				
72	59.396	59.356	58.597	1.345
100	3.691	3.684	3.482	5.662
150	0.064	0.062	0.042	34.375
Crimdon Dune 33'				
72	64.912	64.891	64.192	1.109
100	23.223	23.193	22.188	4.457
150	1.053	1.027	0.834	20.798

	<u>Weight</u>	<u>Weight After Hand Magnet</u>	<u>Weight After Electro- Magnet</u>	<u>% Weight Loss</u>
Crimdon Dune 5'				
72	60.407	60.382	59.891	0.854
100	30.582	30.563	29.908	2.204
150	1.785	1.766	1.616	9.468
Droman Dune B				
72	42.062	42.055	41.572	1.164
100	18.976	18.967	18.077	4.737
150	2.686	2.676	1.360	49.367
200	0.257	0.255	0.013	94.941
Droman Dune A				
72	43.788	43.781	43.354	0.991
100	20.732	20.727	20.070	3.193
150	2.375	2.368	1.175	50.526
200	0.198	0.195	0.009	95.454
Saltburn River Sand				
72	40.302	40.249	37.260	7.548
100	21.173	21.083	20.007	5.507
150	8.912	8.890	--	--
200	2.103	2.094	1.856	0.427
Thorpe Thewles River Sand (4)				
72	13.965	13.954	13.712	1.812
100	6.042	6.032	5.938	1.721
150	3.826	3.818	3.750	1.986
200	1.615	1.608	1.577	2.353
Saltburn River Sand				
Oct. 1967 Flood				
72	38.143	38.040	35.082	8.025
100	21.451	21.379	20.240	5.645
150	8.478	8.447	7.959	6.122
200	1.926	1.917	1.833	31.776

	<u>Weight</u>	<u>Weight After Hand Magnet</u>	<u>Weight After Electro- Magnet</u>	<u>% Weight Loss</u>
Thorpe Thewles Beck (1)				
72	9.074	9.061	8.814	2.865
100	5.776	5.765	5.664	1.939
150	4.191	4.182	4.091	2.386
200	1.941	1.932	1.837	5.358
Thorpe Thewles Beck (2)				
72	12.848	12.830	12.549	2.327
100	7.126	7.116	6.977	2.091
150	4.916	4.909	4.832	1.709
200	2.188	2.180	2.151	1.691
Billingham Beck (1)				
72	25.490	25.464	25.254	1.428
100	16.325	16.311	16.072	1.550
150	9.908	9.891	9.714	1.958
200	3.452	3.446	3.412	1.159
Billingham Beck 2				
72	8.196	8.185	8.029	2.037
100	3.050	3.045	2.989	2.000
150	1.328	1.324	1.238	6.777
200	0.455	0.455	0.446	1.978
Offshore S.L.S. 10				
72	2.046	2.030	1.954	4.496
100	13.307	13.277	12.932	2.818
150	47.931	47.902	46.769	2.424
200	20.745	20.704	20.283	2.227
Offshore S.L.S. 4				
72	19.397	19.386	19.200	1.016
100	19.974	19.960	19.746	1.141
150	33.719	33.693	33.002	2.126
200	13.040	13.013	12.833	1.587

	<u>Weight</u>	<u>Weight After Hand Magnet</u>	<u>Weight After Electro- Magnet</u>	<u>% Weight Loss</u>
Offshore Loc. 2.				
72	2.174	2.161	2.099	3.450
100	18.114	18.101	17.770	1.899
150	57.856	57.805	56.690	2.015
Offshore S.L.S. 3.				
72	17.924	17.918	17.659	1.478
100	21.590	21.555	21.047	2.515
150	28.413	28.367	27.554	3.058
200	11.124	11.065	10.896	2.050
Offshore Loc. 12				
72	49.629	49.629	48.826	1.618
100	28.804	28.797	27.678	3.909
150	3.118	3.095	2.714	12.957
200	0.127	0.123	0.097	23.622
Offshore R.5.				
72	16.519	16.519	16.273	1.489
100	57.488	57.488	56.766	1.256
150	17.549	17.543	17.093	2.598
200	0.619	0.616	0.592	4.361
Offshore R.1.				
72	9.709	9.709	9.516	1.988
100	65.582	65.582	64.911	1.023
150	20.380	20.380	19.876	2.473
200	0.434	0.430	0.409	5.760
Offshore Loc. 16.				
72	43.284	43.268	42.730	1.280
100	27.240	27.240	26.351	3.263
150	4.120	4.086	3.941	4.345
Offshore S.L.S. 9.				
72	3.082	3.065	2.996	2.790
100	18.661	18.648	18.256	2.170
150	57.585	57.533	55.767	3.157
200	14.251	14.165	13.370	6.182

	<u>Weight</u>	<u>Weight After Hand Magnet</u>	<u>Weight After Electro- Magnet</u>	<u>% Weight Loss</u>
Offshore R.2.				
72	12.302	12.293	12.104	1.609
100	57.979	57.962	55.184	4.821
150	22.331	22.319	21.925	1.818
200	0.889	0.878	0.863	2.925
Offshore R.8.				
72	1.783	1.781	1.723	3.365
100	25.096	25.088	24.708	1.546
150	60.470	60.402	58.126	3.876
200	9.409	9.364	8.864	5.792
Offshore Loc.7.				
72	21.362	21.336	20.987	1.755
100	37.560	37.512	35.863	4.518
150	31.385	31.271	29.424	6.248
200	1.701	1.664	1.543	9.289
Offshore Loc.1.				
72	6.937	6.726	6.453	6.977
100	8.455	8.430	8.151	3.595
150	34.143	34.054	33.371	2.261
200	21.789	21.651	21.124	3.051
Offshore S.L.S.6.				
72	11.379	11.369	10.971	3.585
100	18.288	18.271	17.517	4.216
150	47.719	47.662	46.453	2.653
200	12.492	12.427	12.207	2.281
Offshore Loc.10.				
72	44.807	44.766	44.127	1.518
100	26.821	26.783	25.955	3.229
150	5.800	5.774	5.531	4.638
200	0.288	0.288	0.264	8.333

	<u>Weight</u>	<u>Weight After Hand Magnet</u>	<u>Weight After Electro- Magnet</u>	<u>% Weight Loss</u>
Offshore R.9.				
72	18.350	18.330	18.170	0.981
100	64.021	63.996	63.602	0.654
150	14.255	14.242	13.918	2.364
200	0.275	0.267	0.253	8.000
Droman Offshore A. Site 1.				
72	34.336	34.336	33.837	1.453
100	10.551	10.547	10.202	3.307
150	1.484	1.480	1.330	10.377
200	0.232	0.228	0.202	12.931
Droman Offshore B. Site 1.				
72	36.300	36.300	36.098	0.556
100	15.525	15.525	15.402	0.792
150	2.187	2.187	2.120	3.063
200	0.286	0.285	0.270	5.594
Droman Offshore A. Site 2.				
72	40.927	40.925	40.591	0.820
100	17.552	17.548	17.242	1.766
150	2.705	2.701	2.487	8.059
200	0.310	0.310	0.274	11.612
Droman Offshore B. Site 2.				
72	37.235	37.234	36.888	0.931
100	13.271	13.270	13.018	1.906
150	2.759	2.757	2.646	4.095
200	0.480	0.480	0.449	6.458
Droman Offshore C. Site 2.				
72	37.883	37.883	37.641	0.638
100	14.699	14.698	14.510	1.285
150	3.159	3.157	3.065	2.975
200	0.433	0.431	0.411	5.080

	Weight	Weight After Hand Magnet	Weight After Electro- Magnet	% Weight Loss
Coatham Stob Yellow Sand (1)				
72	29.581	29.547	28.883	2.360
100	29.903	29.846	29.179	2.421
150	18.590	18.564	18.045	2.932
200	5.200	5.188	4.880	6.154
Coatham Stob Yellow Sand 3.				
72	33.048	33.026	32.279	2.327
100	27.777	27.743	27.119	2.369
150	13.569	13.551	13.230	2.498
200	3.227	3.196	3.138	2.758
Coatham Stob 4.				
72	25.219	25.175	24.594	2.478
100	20.168	20.133	19.753	0.173
150	11.861	11.838	11.389	3.979
200	3.209	3.190	3.147	1.932
Coatham Stob 1.				
72	23.945	23.908	23.247	2.915
100	12.490	12.438	12.068	3.379
150	4.914	4.868	4.564	7.122
200	1.452	1.429	1.410	2.892
Coatham Stob Same as 3.				
72	41.154	41.112	39.781	3.336
100	15.714	15.690	15.253	2.934
150	5.181	5.069	4.972	4.034
200	1.045	1.036	0.984	5.837
Coatham Stob. 4 (2)				
72	37.616	37.602	36.414	3.195
100	30.516	30.501	29.704	2.660
150	13.874	13.862	13.491	2.760
200	2.690	2.680	2.614	2.825

	<u>Weight</u>	<u>Weight After Hand Magnet</u>	<u>Weight After Electro- Magnet</u>	<u>% Weight Loss</u>
Coatham Stob 4 (2)				
72	29.626	29.606	28.894	2.471
100	32.204	32.181	31.552	2.024
150	18.705	18.698	18.324	2.037
200	4.242	4.232	4.150	2.169
Coatham Stob 5 (2)				
72	7.953	7.944	7.453	6.287
100	4.174	4.160	3.951	5.007
150	0.669	0.669	0.633	5.381
Coatham Stob 3.				
72	30.928	30.899	30.164	2.470
100	30.351	30.321	29.917	1.430
150	19.460	19.441	18.876	3.001
200	6.189	6.161	5.934	4.120
Brierton Top 3A.				
72	30.060	30.037	29.295	2.545
100	27.080	27.060	26.452	2.319
150	14.227	14.218	13.854	2.622
200	4.074	4.070	3.992	2.013
Brierton 2C.				
72	12.638	12.572	11.727	7.208
100	6.662	6.641	6.395	4.008
150	3.249	3.241	2.827	12.989
200	0.838	0.834	0.811	3.222
Brierton M18 3B				
72	20.260	20.223	19.744	2.547
100	21.019	20.995	20.556	2.203
150	24.211	24.203	23.569	2.652
200	11.320	11.256	11.154	1.466

	<u>Weight</u>	<u>Weight After Hand Magnet</u>	<u>Weight After Electro- Magnet</u>	<u>% Weight Loss</u>
Brierton 3C Bottom				
72	17.481	17.466	17.175	1.750
100	27.782	27.761	27.169	2.206
150	25.136	25.113	24.731	1.611
200	11.759	11.727	11.623	1.156
Brierton 4C above 3C				
72	33.329	33.295	32.420	2.727
100	28.594	28.587	27.983	2.137
150	15.499	15.494	15.289	1.355
200	3.402	3.398	3.359	1.264
Brierton A2				
72	25.116	25.077	24.164	3.790
100	14.115	14.097	13.690	3.011
150	7.500	7.492	7.302	2.640
200	2.518	2.514	2.464	2.144
Brierton C2				
72	41.935	41.310	40.079	4.426
100	19.894	19.854	19.114	3.921
150	5.023	5.014	4.794	4.559
200	1.094	1.094	1.035	5.393
Brierton C5				
72	45.775	45.759	44.411	2.980
100	23.654	23.625	22.824	3.509
150	6.026	6.016	5.785	3.999
200	1.099	1.099	1.075	2.184
Wynyard Basal Sand				
72	3.041	3.038	2.958	2.729
100	21.238	21.221	20.937	1.417
150	33.090	33.052	32.506	1.765
200	13.153	13.136	10.870	17.357

	<u>Weight</u>	<u>Weight After Hand Magnet</u>	<u>Weight After Electro- Magnet</u>	<u>% Weight Loss</u>
Wynyard C.				
72	19.759	19.676	18.495	6.397
100	5.795	5.771	5.544	4.331
150	1.740	1.734	1.687	3.046
200	0.310	0.310	0.295	4.839
Wynyard A.				
72	22.857	22.785	21.749	4.847
100	4.326	4.305	4.118	4.808
150	1.312	1.301	1.269	3.277
200	0.274	0.272	0.267	2.555
Wynyard 3				
72	19.203	19.155	18.166	5.400
100	5.960	5.948	5.732	3.825
150	2.133	2.126	2.077	2.625
200	0.636	0.636	0.621	2.358
Wynyard 2.				
72	18.887	18.824	17.667	6.459
100	5.584	5.565	5.332	4.513
150	1.991	1.984	1.538	22.752
200	0.604	0.603	0.590	2.318
Hurworth 3				
72	54.259	54.233	52.896	2.512
100	31.583	31.565	30.450	3.587
150	5.384	5.374	5.115	4.996
200	0.614	0.612	0.572	6.840
Hurworth H3				
72	0.356	0.351	0.335	5.899
100	14.052	14.034	13.700	2.505
150	48.625	48.597	47.802	1.692
200	21.703	21.681	21.367	1.548

	<u>Weight</u>	<u>Weight After Hand Magnet</u>	<u>Weight After Electro- Magnet</u>	<u>% Weight Loss</u>
Hurworth 1				
72	56.685	56.662	54.698	3.505
100	25.107	25.074	23.164	7.739
150	4.306	4.289	4.009	6.897
200	0.556	0.556	0.522	6.115
Hurworth 5				
72	51.631	51.614	50.163	2.843
100	33.415	33.407	31.997	4.244
150	6.162	6.154	6.021	2.288
200	0.873	0.873	0.834	4.467
Hurworth H6				
72	0.255	0.222	0.215	4.444
100	11.846	11.845	11.659	1.578
150	42.892	42.874	41.869	2.385
200	22.861	22.837	22.336	2.296
Newton Bewley 2				
72	13.145	13.116	12.711	3.299
100	9.639	9.629	9.460	1.857
150	10.968	10.948	10.892	0.693
200	5.618	5.581	5.557	1.086
Girsby Flat				
72	15.581	15.549	15.301	1.797
100	19.390	19.390	19.115	1.418
150	17.813	17.813	17.714	0.556
200	8.328	8.324	8.238	1.081
Eaglescliffe 3				
72	1.841	1.833	1.530	16.893
100	9.832	9.827	9.593	2.431
150	44.400	44.398	43.846	1.248
200	23.268	23.250	22.854	1.779

	Weight	Weight After Hand Magnet	Weight After Electro- Magnet	% Weight Loss
Eaglescliffe 4				
72	1.882	1.872	1.804	4.144
100	23.846	23.839	22.678	4.898
150	56.412	56.398	55.426	1.748
200	12.641	12.630	11.800	6.653
Rockcliffe				
72	27.072	27.071	26.570	1.854
100	46.709	46.705	45.942	1.642
150	13.686	13.674	12.535	8.410
200	3.267	3.256	3.137	3.979
Stockton-Norton 1				
72	12.409	12.400	12.023	3.111
100	4.555	4.547	4.374	3.974
150	2.091	2.084	1.994	4.639
200	1.037	1.034	1.005	0.289
Ireland Beach				
72	21.369	21.330	20.995	1.750
100	24.317	24.303	23.605	2.928
150	23.854	23.844	23.202	2.733
200	10.311	10.298	10.197	1.106
Edinburgh Raised Sand				
72	3.949	3.938	3.748	5.090
100	34.439	34.354	33.224	3.527
150	43.647	43.537	42.221	3.267
200	12.276	12.167	11.533	6.052
Edinburgh Raised Sand				
72	3.414	3.405	3.273	4.130
100	36.275	36.083	35.206	2.947
150	46.379	46.282	45.104	2.749
200	10.229	10.138	9.493	7.195

	Weight	Weight After Hand Magnet	Weight After Electro- Magnet	% Weight Loss
<u>Edinburgh Raised Sand</u>				
72	1.620	1.609	1.547	4.506
100	29.627	29.566	28.849	2.626
150	48.607	48.533	47.305	2.679
200	13.894	13.834	13.480	2.980
<u>Edinburgh Fluvio-Glacial</u>				
72	20.457	20.397	19.053	6.863
100	38.272	38.223	36.364	4.985
150	28.437	28.067	26.595	6.477
200	7.346	7.198	6.655	9.406

Other Sedimentary Data

	<u>Micra</u> <u>(grains</u> <u>/1000)</u>	<u>Rock</u> <u>Frag-</u> <u>ments</u> <u>(grains</u> <u>/1000)</u>	<u>Coal</u> <u>(grains</u> <u>/1000)</u>	<u>Shell</u> <u>(grains</u> <u>/1000)</u>	<u>Surface</u> <u>Texture</u>
<u>Seaton Carew Beach 2</u>					
B.S. Standard Sieves					
72	0.0	15	1	138	18
100	0.0	2	1	140	
150	0.0	2	2	120	
200	trace	0.0	34	60	
Saltburn Beach					
72	0.0	28	10	182	19
100	0.0	19	10	170	
150	0.0	11	20	400	
200	0.0	0.0	0.0	79	
Beach North of Saltburn					
72	0.0	32	0.0	103	20
100	0.0	31	0.0	79	
150	0.0	28	10	68	
200	0.0				
Marske Beach (1)					
72	0.0	41	0.0	303	17
100	0.0	5	0.0	130	
150	1.0	4	0.0	114	
200					
South Gare 10					
72	0.0	4	0.0	310	18
100	3	3	0.0	203	
150	0.0	2	0.0	123	
200					

	<u>Micra</u> <u>(grains</u> <u>/1000)</u>	<u>Rock</u> <u>Frag-</u> <u>ments</u> <u>(grains</u> <u>/1000)</u>	<u>Coal</u> <u>(grains</u> <u>/1000)</u>	<u>Shell</u> <u>(grains</u> <u>/1000)</u>	<u>Surface</u> <u>Texture</u>
Redcar Beach (3)					
72	0.0	38	10	413	19
100	0.0	20	8	158	
150	1	4	2	123	
200	0.0	3	2	98	
Musselborough Beach					
72	0.0	83	3	242	17
100	0.0	62	trace	149	
150	0.0	42	0.0	82	
H.Y.C. Beach (3)					
72	0.0	15	40	220	
100	1	10	9	19	
150	8	0.0	7	3	
H.Y.C. Beach (1)					
72	trace	22	110	610	16
100	trace	12	26	441	
150	20	3	39	101	
Crimdon Beach (1)					
72	0.0	0.0	7	30	19
100	0.0	0.0	10	20	
150	0.0	0.0	11	19	
Crimdon Beach (3)					
72	trace	5	10	40	17
100	trace	2	53	41	
150	trace	2	79	40	
Droman Beach A.					
72	1	8	0.0	908	18
100	2	3	0.0	878	
150	0.0	3	0.0	803	
200	-	-	-	-	

	<u>Micra</u> <u>(grains</u> <u>/1000)</u>	<u>Rock</u> <u>Frag-</u> <u>ments</u> <u>(grains</u> <u>/1000)</u>	<u>Coal</u> <u>(grains</u> <u>/1000)</u>	<u>Shell</u> <u>(grains</u> <u>/1000)</u>	<u>Surface</u> <u>Texture</u>
Droman Beach B.					
72	3	2	0.0	623	20
100	2	1	0.0	568	
150	0.0	2	0.0	610	
200	-	-	-	-	
S. Gare Dune D.					
72	0.0	5	0.0	155	13
100	0.0	8	1	89	
150	0.0	2	0.0	152	
S. Gare Dune A.					
72	0.0	4	0.0	84	10
100	0.0	32	4	121	
150	0.0	2	0.0	89	
Cliff Slip Huntcliff					
72	0.0	2	1	7	8
100	0.0	10	0.0	7	
150	0.0	3	20	20	
Musselborough Sand Dune					
72	0.0	72	10	101	13
100	0.0	50	2	109	
150	0.0	4	1	111	
200	0.0	2	2	45	
Crimdon Dune 2					
72	0.0	18	0.0	32	12
100	0.0	18	15	38	
150	0.0	24	25	56	
Crimdon Dune 18'					
72	0.0	3	0.0	3	11
100	0.0	21	12	98	
150	0.0	23	2	82	

	<u>Micra</u> <u>(grains</u> <u>/1000)</u>	<u>Rock</u> <u>Frag-</u> <u>ments</u> <u>(grains</u> <u>/1000)</u>	<u>Coal</u> <u>(grains</u> <u>/1000)</u>	<u>Shell</u> <u>(grains</u> <u>/1000)</u>	<u>Surface</u> <u>Texture</u>
Crimdon Dune 33'					
72	0.0	5	0.0	110	14
100	0.0	0.0	3	80	
150	0.0	10	20	110	
Crimdon Dune 5'					
72	0.0	10	0.0	101	12
100	0.0	20	2	70	
150	0.0	15	15	48	
Droman Dune B.					
72	0.0	3	0.0	586	14
100	0.0	2	0.0	632	
150	0.0	0.0	0.0	203	
200	0.0	4	0.0	58	
Droman Dune A					
72	0.0	1	0.0	678	11
100	0.0	5	0.0	312	
150	0.0	3	0.0	88	
200	0.0	2	0.0	59	
Saltburn River Sand					
72	0.0	140	Trace	0.0	8
100	8	130	28	0.0	
150	7	40	25	0.0	
200	10	28	29	0.0	
Thorpe Thewles River Sand 4.					
72	1	33	8	0.0	9
100	1	14	9	0.0	
150	4	12	10	0.0	
200	8	0.0	4	0.0	

	<u>Micra</u> (grains /1000)	<u>Rock</u> <u>Frag-</u> <u>ments</u> (grains /1000)	<u>Coal</u> (grains /1000)	<u>Shell</u> (grains /1000)	<u>Surface</u> <u>Texture</u>
Saltburn River Sand					
Oct. 1967 Flood					
72	trace	30	0.0	0.0	9
100	1	11	0.0	0.0	
150	0.0	0.0	0.0	0.0	
200	0.0	0.0	0.0	0.0	
Thorpée Thewles Beck 1.					
72	0.0	530	0.0	0.0	7
100	0.0	361	0.0	0.0	
150	1	239	38	0.0	
200	0.0	219	36	0.0	
Thorpée Thewles Beck 2.					
72	0.0	90	0.0	trace	10
100	0.0	50	0.0	trace	
150	0.0	26	5	0.0	
200	0.0	8	27	0.0	
Billingham Beck (1)					
72	2	130	10	0.0	9
100	0.0	30	2	0.0	
150	2	34	2	0.0	
200	1	40	11	0.0	
Billingham Beck 2.					
72	0.0	40	0.0	0.0	8
100	0.0	10	0.0	0.0	
150	6	65	21	0.0	
200	2	73	18	0.0	
Offshore S.L.S.10					
72	22	20	434	251	13
100	2	7	272	107	
150	1	3	181	84	
200	2	0.0	140	61	

	<u>Micra</u> <u>(grains</u> <u>/1000)</u>	<u>Rock</u> <u>Frag-</u> <u>ments</u> <u>(grains</u> <u>/1000)</u>	<u>Coal</u> <u>(grains</u> <u>/1000)</u>	<u>Shell</u> <u>(grains</u> <u>/1000)</u>	<u>Surface</u> <u>Texture</u>
Offshore S.L.S. 4					
72	0.0	30	10	168	14
100	0.0	31	8	111	
150	3	13	7	71	
200	9	2	173	4	
Offshore LOC 2.					
72	10	0.0	513	335	12
100	1	38	270	104	
150	19	0.0	218	114	
Offshore S.L.S.3					
72	0.0	11	62	359	14
100	0.0	3	20	203	
150	3	4	85	110	
200	14	4	90	18	
Offshore LOC.12					
72	2	11	7	160	15
100	8	11	13	90	
150	3	8	43	92	
200	5	23	80	41	
Offshore R.5					
72	0.0	10	10	362	9
100	0.0	7	8	158	
150	trace	3	4	92	
200	trace	trace	3	78	
Offshore R.1					
72	5	8	12	303	12
100	2	trace	2	212	
150	trace	4	2	83	
200	3	0.0	23	91	
Offshore LOC.16					
72	0.0	40	31	310	14
100	0.0	10	30	150	
150	0.0	3	45	111	

	<u>Micra</u> <u>(grains</u> <u>/1000)</u>	<u>Rock</u> <u>Frag-</u> <u>ments</u> <u>(grains</u> <u>/1000)</u>	<u>Coal</u> <u>(grains</u> <u>/1000)</u>	<u>Shell</u> <u>(grains</u> <u>/1000)</u>	<u>Surface</u> <u>Texture</u>
Offshore S.L.S.9					
72	0.0	37	142	264	12
100	0.0	36	15	111	
150	2	20	11	61	
200	1	18	49	2	
Offshore R.2					
72	0.0	10	0.0	170	8
100	0.0	0.0	0.0	10	
150	0.0	19	17	0.0	
200	0.0	7	2	3	
Offshore R.8					
72	0.0	3	11	372	11
100	1	5	7	103	
150	3	0.0	5	58	
200	5	0.0	2	20	
Offshore LOC.7					
72	0.0	29	21	60	13
100	0.0	19	50	68	
150	0.0	22	41	32	
200	trace	12	38	28	
Offshore LOC 1					
72	12	32	177	165	13
100	7	23	434	89	
150	70	15	111	48	
200	3	25	72	23	
Offshore S.L.S.6					
72	9	43	50	210	13
100	1	12	45	54	
150	trace	11	29	10	
200	trace	8	10	10	
Offshore LOC.10					
72	0.0	8	32	34	12
100	0.0	19	18	140	
150	trace	38	31	17	
200	0.0	12	4	51	

	<u>Micra</u> (grains /1000)	<u>Rock</u> <u>Frag-</u> <u>ments</u> (grains /1000)	<u>Coal</u> (grains /1000)	<u>Shell</u> (grains /1000)	<u>Surface</u> <u>Texture</u>
Offshore R.9					
72	trace	1	1	300	10
100	trace	0.0	1	51	
150	trace	0.0	3	30	
200	trace	0.0	5	11	
Droman Offshore A Site 1.					
72	5	9	0.0	450	12
100	4	48	0.0	500	
150	3	28	0.0	410	
200	4	12	0.0	420	
Droman Offshore B.Site 1					
72	8	25	0.0	580	14
100	0.0	12	0.0	420	
150	1	19	0.0	520	
200	0.0	135	0.0	350	
Droman Offshore A.Site 2.					
72	0.0	80	0.0	400	15
100	0.0	55	0.0	425	
150	0.0	36	0.0	350	
200	0.0	142	0.0	284	
Droman Offshore B.Site 2.					
72	0.0	11	0.0	743	11
100	0.0	4	0.0	648	
150	0.0	5	0.0	724	
200	0.0	35	0.0	803	
Droman Offshore C. Site 2.					
72	2	17	0.0	621	10
100	1	13	0.0	785	
150	0.0	15	0.0	652	
200	0.0	65	0.0	429	
Coatham Stob Yellow Sand 1.					
72	0.0	41	8	0.0	11
100	0.0	10	3	0.0	
150	5	50	0.0	0.0	
200	12	15	0.0	trace	

	<u>Micra</u> <u>(grains</u> <u>/1000)</u>	<u>Rock</u> <u>Frag-</u> <u>ments</u> <u>(grains</u> <u>/1000)</u>	<u>Coal</u> <u>(grains</u> <u>/1000)</u>	<u>Shell</u> <u>(grains</u> <u>/1000)</u>	<u>Surface</u> <u>Texture</u>
Coatham Stob Yellow Sand 3.					
72	trace	23	2	0.0	7
100	0.0	30	31	0.0	
150	trace	20	4	0.0	
200	trace	13	10	0.0	
Coatham Stob 4					
72	10	83	23	0.0	14
100	10	41	17	0.0	
150	11	18	trace	0.0	
200	10	0.0	4	0.0	
Coatham Stob 1.					
72	0.0	39	2	0.0	11
100	0.0	21	38	0.0	
150	0.0	17	57	0.0	
200	5	0.0	78	0.0	
Coatham Stob, Same as 3.					
72	0.0	14	11	0.0	8
100	0.0	5	2	0.0	
150	0.0	3	0.0	0.0	
200	31	0.0	5	0.0	
Coatham Stob 4.(2)					
72	0.0	54	22	trace	10
100	0.0	23	8	0.0	
150	0.0	15	2	0.0	
200	0.0	11	1	trace	
Coatham Stob 4(1)					
72	1	37	22	trace	9
100	3	32	20	trace	
150	10	33	20	trace	
200	11	3	21	trace	
Coatham Stob 5(2)					
72	0.0	70	32	trace	12
100	trace	42	25	trace	
150	1	41	10	trace	

	<u>Micra</u> <u>(grains</u> <u>/1000)</u>	<u>Rock</u> <u>Frag-</u> <u>ments</u> <u>(grains</u> <u>/1000)</u>	<u>Coal</u> <u>(grains</u> <u>/1000)</u>	<u>Shell</u> <u>(grains</u> <u>/1000)</u>	<u>Surface</u> <u>Texture</u>
Coatham Stob 3					
72	trace	18	10	0.0	11
100	trace	15	7	0.0	
150	1	2	27	0.0	
200	1	2	1	0.0	
Brierton, Top 3A					
72	0.0	165	3	0.0	7
100	0.0	45	15	0.0	
150	2	3	9	0.0	
200	5	2	12	0.0	
Brierton 2C					
72	0.0	130	11	0.0	6
100	1	42	12	0.0	
150	7	20	10	0.0	
200	5	0.0	10	0.0	
Brierton Mid 3B					
72	1	52	40	0.0	10
100	20	30	21	0.0	
150	8	19	11	0.0	
200	11	3	6	0.0	
Brierton 3C.Bottom					
72	0.0	61	27	0.0	8
100	4	40	30	0.0	
150	9	12	20	0.0	
200	8	2	10	0.0	
Brierton 4C above 3C					
72	1	71	28	0.0	9
100	2	22	10	0.0	
150	1	11	8	0.0	
200	trace	2	8	0.0	
Brierton A2					
72	trace	28	0.0	0.0	8
100	10	28	trace	0.0	
150	9	7	1	0.0	
200	11	0.0	2	0.0	

	<u>Micra</u> (grains /1000)	<u>Rock</u> <u>Frag-</u> <u>ments</u> (grains /1000)	<u>Coal</u> (grains /1000)	<u>Shell</u> (grains /1000)	<u>Surface</u> <u>Texture</u>
Brierton C2					
72	10	28	10	0.0	8
100	0.0	31	11	0.0	
150	3	30	8	0.0	
200	5	5	3	0.0	
Brierton C5					
72	0.0	62	trace	0.0	7
100	0.0	54	trace	0.0	
150	1	41	trace	0.0	
200	5	13	trace	0.0	
Wynyrd Basal Sand					
72	2	332	10	0.0	13
100	1	96	3	0.0	
150	1	72	1	0.0	
200	2	60	20	0.0	
Wynyrd C.					
72	trace	240	trace	0.0	6
100	trace	180	0.0	0.0	
150	trace	150	trace	0.0	
200	trace	160	trace	0.0	
Wynyrd A.					
72	0.0	168	0.0	0.0	7
100	0.0	99	trace	0.0	
150	0.0	71	0.0	0.0	
200	1	41	0.0	0.0	
Wynyrd 3.					
72	0.0	200	0.0	0.0	8
100	0.0	309	0.0	0.0	
150	0.0	260	trace	0.0	
200	0.0	260	trace	0.0	
Wynyrd 2.					
72	0.0	180	0.0	0.0	7
100	0.0	130	5	0.0	
150	trace	103	0.0	0.0	
200	trace	71	0.0	0.0	

	<u>Micra</u> <u>(grains</u> <u>/1000)</u>	<u>Rock</u> <u>Frag-</u> <u>ments</u> <u>(grains</u> <u>/1000)</u>	<u>Coal</u> <u>(grains</u> <u>/1000)</u>	<u>Shell</u> <u>(grains</u> <u>/1000)</u>	<u>Surface</u> <u>Texture</u>
Hurworth 3					
72	trace	98	100	0.0	10
100	trace	10	20	0.0	
150	trace	10	15	0.0	
200	1	9	11	0.0	
Hurworth H.3					
72	111	233	183	0.0	11
100	84	70	53	0.0	
150	2	25	32	0.0	
200	2	0.0	10	0.0	
Hurworth 1.					
72	0.0	90	20	0.0	10
100	0.0	60	30	0.0	
150	0.0	170	11	0.0	
200	0.0	41	17	0.0	
Hurworth 5.					
72	0.0	39	11	0.0	12
100	0.0	31	10	0.0	
150	0.0	29	19	0.0	
200	0.0	42	23	0.0	
Hurworth H6					
72	106	100	75	0.0	14
100	29	30	45	0.0	
150	9	0.0	15	0.0	
200	2	0.0	15	0.0	
Newton Bewley 2.					
72	trace	250	0.0	0.0	8
100	trace	58	trace	0.0	
150	2	80	0.0	0.0	
200	2	8	0.0	0.0	
Girsby Flat					
72	234	521	178	0.0	10
100	92	183	42	0.0	
150	41	72	19	0.0	
200	10	20	2	0.0	

	<u>Micra</u> <u>(grains</u> <u>/1000)</u>	<u>Rock</u> <u>Frag-</u> <u>ments</u> <u>(grains</u> <u>/1000)</u>	<u>Coal</u> <u>(grains</u> <u>/1000)</u>	<u>Shell</u> <u>(grains</u> <u>/1000)</u>	<u>Surface</u> <u>Texture</u>
Eaglescliffe 3.					
72	50	341	149	0.0	12
100	20	110	28	0.0	
150	2	67	2	0.0	
200	11	6	1	0.0	
Eaglescliffe 4.					
72	70	682	73	0.0	14
100	59	151	20	0.0	
150	27	32	5	0.0	
200	1	2	3	0.0	
Rockcliffe					
72	1	41	30	0.0	11
100	1	24	19	0.0	
150	3	17	21	0.0	
200	1	7	61	0.0	
Stockton-Norton 1.					
72	0.0	18	8	0.0	12
100	2	3	1	0.0	
150	8	0.0	1	0.0	
200	22	0.0	0.0	0.0	
Ireland Beach					
72	0.0	2	0.0	1	17
100	0.0	1	0.0	2	
150	4	0.0	0.0	3	
200	2	0.0	0.0	3	
Edinburgh Raised Sand					
72	300	89	310	0.0	9
100	1	112	31	0.0	
150	5	13	50	0.0	
200	1	51	40	0.0	
Edinburgh Raised Sand					
72	70	90	61	0.0	10
100	2	20	15	0.0	
150	7	20	39	0.0	
200	2	4	3	0.0	

	<u>Micra</u> <u>(grains</u> <u>/1000)</u>	<u>Rock</u> <u>Frag-</u> <u>ments</u> <u>(grains</u> <u>/1000)</u>	<u>Coal</u> <u>(grains</u> <u>/1000)</u>	<u>Shell</u> <u>(grains</u> <u>/1000)</u>	<u>Surface</u> <u>Texture</u>
Edinburgh Raised Sand					
72	62	89	120	0.0	8
100	23	41	42	0.0	
150	3	52	60	0.0	
200	3	51	50	0.0	
Edinburgh Fluvial-glacial.					
72	5	72	72	0.0	11
100	1	40	18	0.0	
150	2	15	10	0.0	
200	3	4	7	0.0	

