



Durham E-Theses

Ecology of ragadia makuta (lepidoptera: satyrinae) in tropical rainforests of Sabah, Malaysia.

Dawood, Mahadimenakbar Mohamed

How to cite:

Dawood, Mahadimenakbar Mohamed (2000) *Ecology of ragadia makuta (lepidoptera: satyrinae) in tropical rainforests of Sabah, Malaysia.*, Durham theses, Durham University. Available at Durham E-Theses Online: <http://etheses.dur.ac.uk/938/>

Use policy

The full-text may be used and/or reproduced, and given to third parties in any format or medium, without prior permission or charge, for personal research or study, educational, or not-for-profit purposes provided that:

- a full bibliographic reference is made to the original source
- a [link](#) is made to the metadata record in Durham E-Theses
- the full-text is not changed in any way

The full-text must not be sold in any format or medium without the formal permission of the copyright holders.

Please consult the [full Durham E-Theses policy](#) for further details.

is confined in closed-canopy forest (Hill 1999). In addition, the food plants for this butterfly (*Selagenella* spp.) can be readily recorded from ground-based survey making it possible to quantify habitat requirements (Hill 1999).



Figure 1.3: *R.makuta* perching on *Selagenella* sp.

This thesis investigates the distribution and abundance of *R.makuta* in unlogged and logged forest selectively logged 10 to 11 years previously. Chapter two describes the study site and general materials and methods used to survey butterflies. Chapter three investigates habitat requirements of *R.makuta* and the effects of habitat modification on distribution and abundance of *R.makuta*. Chapter three also describes the methods used to measure and analyze vegetation structure at the study sites. Chapter four investigates seasonal effects of rainfall and sunshine on *R.makuta* abundance and chapter five discusses the use of *R.makuta* as a bioindicator for other forest butterflies.

analysis. Two people took readings at every station and the mean value was used in the analysis.



Figure 3.1: Densiometer used to measure canopy openness.

Abundance of *Selagenella* spp. was recorded at each station because this herb is the larval food-plant of *Ragadia makuta*. Abundance of *Selagenella* spp. was measured along 10 m transects running north, south, east and west from the centre of the station. Each transect was divided into 20 cm sections using a tape measure and the number of sections (200 per station) where a *Selagenella* plant touched the tape was recorded. The distance of each station to the nearest stream was also recorded because this is likely to determine the probability of *R.makuta* presence. Meteorological data (temperature, monthly rainfall and sunshine) were obtained from the meteorological station at the field centre.

Vegetation variables measured in this study are likely to be closely correlated and were analysed by principle component analysis (PCA). PCA is a data reduction technique used to reduce a large number of closely related variables to a smaller set of